



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

UC-NRLF



5B 50 845

YC 40029

LIBRARY

OF THE

University of California.

No.

14008


Division

Range


Shelf


Received

March 1879










Clarendon Press Series

A TREATISE
ON THE
KINETIC THEORY OF GASES .

BY

HENRY WILLIAM WATSON, M.A.

FORMERLY FELLOW OF TRINITY COLLEGE, CAMBRIDGE



Oxford

AT THE CLARENDON PRESS

1876

1175
W3

London

MACMILLAN AND CO.



14008

PUBLISHERS TO THE UNIVERSITY OF

Oxford

115

PREFACE.

THE idea of a Kinetic Theory of Gases originated with J. Bernouilli about the middle of the last century, but the first establishment of the theory on a scientific basis is due to Professor Clausius.

During the last few years the theory has been greatly developed by many physicists, especially by Professor Clerk Maxwell in England and Professor Clausius and Dr. Ludwig Boltzmann on the Continent; and although still beset by formidable difficulties, it has succeeded in explaining most of the established laws of gases in so remarkable a manner as to render it well worthy of the attentive consideration of scientific men.

My great object in the following short treatise is to make the existing state of the theory more widely known by presenting some of the scattered memoirs of the writers I have mentioned in a systematic and continuous form, in the hope that mathematicians may be induced to turn their attention to the theory, and thus assist in removing, if possible, the obstacles which yet remain in the way of its complete establishment.

For the most part I have followed the method of treatment adopted by Dr. Ludwig Boltzmann in some very interesting memoirs contributed by him to the

Transactions of the Imperial Academy of Vienna,* but in some cases I have varied this treatment for the sake of greater conciseness or greater generality.

* Thus, in place of Dr. L. Boltzmann's conception of a molecule as a collection of mutually attracting particles, I have substituted the more general conception of a material system possessing a given number of degrees of freedom, that is to say, a given number of generalised coordinates.

Again, in the deduction of the second law of Thermodynamics from the results of the Kinetic Theory, I felt some difficulty in following Dr. Boltzmann's reasoning, and I originally proposed to substitute for it a demonstration of my own, free from what appeared to me to be the obscurities of Dr. Boltzmann's reasoning, but applicable only to the case in which there were no intermolecular actions. My friend Mr. S. H. Burbury, formerly fellow of St. John's College, Cambridge, to whom I communicated my difficulties, has invented an unexceptionable proof applicable to all cases, which he published last January in the London, Edinburgh, and Dublin Philosophical Magazine, and with his permission I have adopted this proof in the following treatise.

To Professor Clerk Maxwell I am indebted for much kind assistance, and especially for access to some of his manuscript notes on this subject, from which I have taken many valuable suggestions.

H. W. WATSON.

BERKSWELL RECTORY,

COVENTRY, *Sept. 17, 1876.*

* *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften* Wien, Band 63, 1871, und Band 66, 1872.

LIBRARY
UNIVERSITY OF
CALIFORNIA.

INTRODUCTION.

THE Kinetic Theory of Gases is based upon the conception of an infinitely large number of molecules in motion in a given space with velocities of all degrees of intensity and in all conceivable directions. These molecules, as will be explained in the course of the following treatise, may sometimes be regarded as smooth spheres, in which case we shall only have to consider the motion of translation of the centre of mass of each of them, or they may be regarded as bodies of any form capable of any number of internal vibrations. It is clear that the individual molecules in such a system must be continually acting upon each other, either in the way of collision, like the mutual impacts of elastic spheres, or else in the more gradual way of mutual attraction and repulsion; such actions are called encounters.

It is easy to see that if encounters take place among a great number of molecules, their velocities, even if originally equal, will become unequal, for, except under conditions which can be only rarely satisfied, two molecules having equal velocities before their encounter will have unequal velocities after such encounter.

Now, as long as we have to deal with only two molecules, and have all the data of an encounter given us, we can calculate the result of their mutual action; but when we have to deal with millions of molecules, each of which has millions of encounters in a second, the complexity of the problem seems to shut out all hope of a legitimate solution.

We are obliged therefore to abandon the strictly kinetic method and to adopt the statistical method.

According to the strict kinetic or historical method as applied to the case before us, we follow the whole course of every individual molecule. We arrange our symbols so as to be able to identify every molecule throughout its motion, and the complete solution of the problem would enable us to determine at any given instant the position and motion of any given molecule from a knowledge of the positions and motions of all the molecules in their initial state.

✓ According to the statistical method, the state of the system at any instant is ascertained by distributing the molecules into groups, the definition of each group being founded on some variable property of the molecules. Each individual molecule is sometimes in one of these groups and sometimes in another, but we make no attempt to follow it; we simply take account of the number of molecules which at a given instant belong to each group.

Thus we may consider as a group those molecules which at a given instant lie within a given region of space. Molecules may pass into or out of this region, but we confine our attention to the increase or diminution of the number of molecules within it. Just as the population of a watering-place, considered as a mere number, varies in the same way whether its visitors return to it season after season, or whether the annual gathering consists each year of fresh individuals. Or we may form our group out of those molecules which at a given instant have velocities lying within given limits. When a molecule has an encounter and changes its velocity, it passes out of one of these groups and enters another; but as other molecules are also changing their velocities, the number of molecules in each group varies little from a certain average value.

✓ We thus meet with a new kind of regularity, the regularity of averages, a regularity which, when we are dealing with millions of millions of individuals, is so unvarying that we

are almost in danger of confounding it with absolute uniformity.*

Laplace, in his Theory of Probability, has given many examples of this kind of statistical regularity, and has shown how this regularity is consistent with the utmost irregularity among the individual instances which are enumerated in making up the results.*

These observations must be borne in mind in interpreting the definitions laid down and the results arrived at in the Kinetic Theory of Gases.

Thus, to refer to the illustrations already given, we shall prove that the number of molecules lying within a certain region of space, or the number of molecules having their velocities within certain limits differing by some finite quantity, is in each case a number bearing some finite ratio to the total number of molecules in the mass under consideration, and therefore infinitely large. But these results are to be interpreted as average results. We do not assert by them, nor are we capable of proving, that at any given instant there is one single molecule satisfying either of the required conditions, that is, comprised within either of the contemplated groups.

So, again, the density of the region in the neighbourhood of any point is defined as the limit of the quotient of the number representing the aggregate masses of the molecules within any volume containing the point, to the number representing that volume, when the volume is indefinitely diminished. In interpreting this definition two things must be remembered.* In the first place, according to what has been said just now, we do not assert and cannot prove that there is, as a matter of fact, any particular number of molecules within the volume containing the given point, at any given instant; and in the second place, supposing we could prove that the number of molecules within the volume was thus accurately determined, yet even so there could be no

* MS. notes by Professor Clerk Maxwell.

point within the region at which the actual density of the matter had the value determined by our definition; for if the point were within a molecule the actual density would be much greater, and if it were not within a molecule the density would be zero.

LIBRARY UNIVERSITY OF CALIFORNIA. KINETIC ~~THEORY~~ OF GASES.

PROPOSITION I.

A VERY great number of smooth elastic spheres equal in every respect are in motion within a region of space of given volume, and therefore occasionally impinge upon each other with various degrees of relative velocity and in various directions. The space is so large in proportion to the sum of the volumes of the spheres that the average time during which any one sphere is moving free from contact with any other is infinitely greater than the average time during which it is in collision with some other sphere*. Required to find the law according to which the velocities must be distributed in order that such distribution may be permanent.

Let N be the total number of spheres, and let

$$\chi(u, v, w) du dv dw$$

be the number of spheres whose component velocities parallel to the axes are intermediate between

$$u \text{ and } u + du, \quad v \text{ and } v + dv, \quad w \text{ and } w + dw$$

respectively.

If we change the variables and make c the resultant velocity, θ the inclination of c to the axis of z , and ϕ that of the plane of cz to the plane of xz , the expression given above will become

$$\chi(u, v, w) c^2 \sin \theta d\theta d\phi dc.$$

Let a spherical surface of radius unity be described round the origin as centre, and let us write $d\sigma$ for the element $\sin \theta d\theta d\phi$ on this spherical surface, then the last written expression becomes

$$\chi(u, v, w) c^2 dc d\sigma.$$

* In the mathematical conception of a collision as an absolutely instantaneous phenomenon the proviso in the text is of course superfluous. An actual physical collision must take some time, however short, and the object of the proviso is to exclude the possibility, or rather to diminish indefinitely the probability, of the occurrence of cases in which the collision of one sphere with a second is not concluded before that with a third commences, and also to ensure that the number of colliding molecules at any instant is very much less than the number of molecules free from collision.

✓ Since for the same magnitude of the resultant velocity all directions must be equally probable, it follows that the coefficient of $dc d\sigma$ in this expression must be a function of the resultant velocity c only, and therefore the number of spheres having component velocities between u and $u + du$, v and $v + dv$, w and $w + dw$ must be

$$\psi(c) du dv dw.$$

It is required, now, to find the form of ψ in order that the value of this expression may be unaffected by collisions.

We assume that in the permanent state the distribution of the spheres throughout the space occupied by them is homogeneous in all respects; that is to say, on an average of any long time there are the same number of spheres in a given volume wherever that volume may be situated, and the law of distribution of velocities is the same throughout that volume
* as in the whole region under consideration.

Hence the number of pairs of spheres having component velocities between

$$u_1 \text{ and } u_1 + du_1, \quad v_1 \text{ and } v_1 + dv_1, \quad w_1 \text{ and } w_1 + dw_1$$

for the one, and

$$u_2 \text{ and } u_2 + du_2, \quad v_2 \text{ and } v_2 + dv_2, \quad w_2 \text{ and } w_2 + dw_2$$

for the other, and such that the lengths of the projections of their line of centres upon the axes are between

$$x \text{ and } x + dx, \quad y \text{ and } y + dy, \quad z \text{ and } z + dz,$$

must be proportional to

$$\psi(c_1) \psi(c_2) du_1 dv_1 dw_1 du_2 dv_2 dw_2 dx dy dz, \quad \dots \dots \dots (A)$$

where c_1 and c_2 are the resultant velocities.

Since the coefficient of $dx dy dz$ is independent of x, y, z , it follows that the number of pairs having their velocities between the above-mentioned limits and the projections of their line of centres between x and $x + dx$, 0 and dy , 0 and dz must be also represented by (A).

If dy and dz be infinitesimal in comparison with the diameter of any sphere, the lines of centres are parallel to the axis of x and the velocity of approach of these centres is $u_1 - u_2$.

If d be the sum of the radii of any two of the spheres, or, what is the same thing in this case, the diameter of any one of them, then it follows that any pair of spheres having the x component of their line of centres less than $d + (u_1 - u_2)dt$ will encounter each other in the time dt ; making therefore

$$x = d \text{ and } dx = (u_1 - u_2)dt,$$

it will follow that the number of collisions in time dt of pairs of spheres with their lines of centres parallel to x and velocities between the above limits is proportional to

$$\psi(c_1) \psi(c_2) du_1 \dots dw_2 dy dz (u_1 - u_2) dt. \dots \dots (B)$$

After collision let the different quantities involved in (B) become c'_1, c'_2, u'_1 , etc., while x, y , and z, dy and dz remain of course unchanged.

It follows therefore that if the values of these velocities and component velocities had originally been within the limits indicated by the accented letters with reversed signs, the spheres would, after collision, have passed into the original state as indicated by the unaccented letters. Call these states E and F respectively. For permanence of distribution therefore it is necessary that the number of collisions of pairs of spheres in state E during the time dt should be equal to the number of collisions of pairs in the state F during the same time; and therefore that

$$\begin{aligned} \psi(c_1) \psi(c_2) du_1 \dots dw_2 dy dz (u_1 - u_2) \\ = \psi(c'_1) \psi(c'_2) du'_1 \dots dw'_2 dy dz (u'_2 - u'_1). \end{aligned}$$

Now	$u'_1 = u_2,$	$\therefore du'_1 = du_2,$
	$u'_2 = u_1,$	$du'_2 = du_1,$
	$v_1 = v'_1,$	$dv_1 = dv'_1,$
	$v_2 = v'_2,$	$dv_2 = dv'_2,$
	$w = w'_1,$	$dw_1 = dw'_1,$
	$w_2 = w'_2,$	$dw_2 = dw'_2.$

Hence

$$du_1 \dots dw_2 dy dz (u_1 - u_2) = du'_1 \dots dw'_2 dy dz (u'_2 - u'_1);$$

$$\therefore \psi(c_1) \psi(c_2) = \psi(c'_1) \psi(c'_2), \quad \checkmark$$

and

$$c_1^2 + c_2^2 = c_1'^2 + c_2'^2.$$

This functional equation may be integrated as follows. Let it be expressed in the form

$$\chi(c_1^2) \chi(c_2^2) = \chi(c_1'^2) \chi(c_2'^2),$$

and for c_1^2 and c_2^2 write x and x' , and for $c_1'^2$ and $c_2'^2$ write y and y' ; then we have

$$\chi(x)\chi(x') = \chi(y)\chi(y'),$$

where

$$x + x' = y + y';$$

$$\therefore \chi(x)\chi(x') = \chi(y)\chi(x + x' - y);$$

therefore differentiating with respect to x and x' ,

$$\chi'(x)\chi(x') = \chi(y)\chi'(x + x' - y);$$

and

$$\chi(x)\chi'(x') = \chi(y)\chi'(x + x' - y);$$

$$\therefore \frac{\chi'(x)}{\chi(x)} = \frac{\chi'(x')}{\chi(x')};$$

that is,

$$\frac{\chi'(x)}{\chi(x)} = \text{constant};$$

$$\therefore \chi(x) = Ae^{-hx},$$

$$\therefore \psi(c) = Ae^{-hc^2}.$$

Now the axis of x may be taken in any direction. Whence it follows that this form of ψ will ensure the permanence of distribution for all possible collisions.

Therefore the number of spheres with component velocities between u and $u + du$, v and $v + dv$, w and $w + dw$

is

$$Ae^{-hc^2} du dv dw,$$

or

$$Ae^{-hc^2} c^2 dc d\sigma.$$

employing the notation used above.

Integrating with respect to $d\sigma$ from 0 to 4π we find for the number of spheres with velocities between c and $c + dc$ the expression

$$4\pi Ae^{-hc^2} c^2 dc.$$

Again, since the number with component velocities between

$$u \text{ and } u + du, \quad v \text{ and } v + dv, \quad w \text{ and } w + dw$$

is

$$Ae^{-h(u^2+v^2+w^2)} du dv dw,$$

or

$$\sqrt[3]{Ae^{-hu^2}} du \sqrt[3]{Ae^{-hv^2}} dv \sqrt[3]{Ae^{-hw^2}} dw,$$

it follows that the number of spheres having velocities intermediate between u and $u + du$ parallel to any fixed line is

$$Ae^{-hu^2} du \int_{-\infty}^{\infty} e^{-hv^2} dv \int_{-\infty}^{\infty} e^{-hw^2} dw;$$

that is

$$Ae^{-hu^2} du \frac{\pi}{h};$$

where A may be determined by the equation

$$A \frac{\pi}{h} \int_{-\infty}^{\infty} e^{-hu^2} du = N;$$

or

$$A \frac{\pi^{\frac{3}{2}}}{h^{\frac{3}{2}}} = N;$$

and therefore

$$A = \frac{N h^{\frac{3}{2}}}{\pi^{\frac{3}{2}}},$$

that is to say, the number of spheres having velocities between c and $c + dc$ is

$$\frac{4 N h^{\frac{3}{2}}}{\sqrt{\pi}} \cdot e^{-hc^2} c^2 dc.$$

$$h = \frac{1}{2} \frac{m}{\pi t}$$

If we multiply this expression by c , integrate the product with regard to c from 0 to ∞ and divide by N , we find the mean velocity of all the spheres to be

$$\frac{2}{\sqrt{\pi h}}. \quad *$$

And, similarly, multiplying by c^2 instead of c , we find the mean square of the velocities to be

$$\frac{3}{2h}, \quad *$$

and this is greater than the square of the mean velocity as it ought to be.

Before we proceed further it will be useful to make a formal statement of the method employed in the last proposition for denoting the velocities of molecules or material points. ✓

Instead of drawing straight lines from each molecule indicating the magnitude and direction of the velocity of that molecule, we draw all such straight lines for all molecules from any assumed point taken as the origin.

This method is very useful when, as in investigations like the present, we wish to compare the simultaneous velocities of different molecules as well as the successive velocities of each molecule separately.

We thus obtain a figure every point of which corresponds to one of our molecules, the velocity of that molecule is represented in magnitude and direction by a line drawn from the origin to the corresponding point, and the relative velocity of any two

✓ molecules is also represented by the line joining the points corresponding to these two molecules. If the system have a common velocity, then we must suppose the position of the new origin of the diagram of velocities with respect to the old origin to be so chosen as to represent this velocity.

In studying the motion of the system it is found convenient to divide the molecules into groups according to their velocities, those molecules whose velocities lie within certain limits with respect to magnitude and direction being placed in the same group.

In the velocity diagram these molecules are indicated at once by the points which correspond to them being included within a certain small region or elementary volume of the diagram, the boundary of this region corresponding to the given limits of velocity.

Thus, in the proposition just now considered, the state E might be described as that in which the velocity points are situated within the elementary volume

$$du dv dw \text{ or } c^2 \sin \theta d\theta d\phi dc$$

of the diagram of velocities.

We may also conveniently make use of the term velocity-density to indicate the result of dividing the number of molecules whose velocities lie within the given limits by the volume of the corresponding region in the diagram of velocities.

PROPOSITION II.

Let the N elastic spheres of the last proposition be replaced by two sets, one of N spheres each having mass m , and the other of N' spheres each having mass m' , and let us find the law of distribution of the velocities of the N and N' spheres respectively.

Exactly as in the last proposition, it may be proved that the number of N spheres whose component velocities parallel to the axes are intermediate between

u and $u+du$, v and $v+dv$, w and $w+dw$ respectively must be

$$\psi(c) du dv dw,$$

where

$$c^2 = u^2 + v^2 + w^2;$$

and that the number of the N' spheres whose component velocities are intermediate between

$$u' \text{ and } u' + du', \quad v' \text{ and } v' + dv', \quad w' \text{ and } w' + dw'$$

must be

$$\psi_1(c') du' dv' dw';$$

and that the number of collisions in time dt between pairs of spheres, one from each set, with lines of centres parallel to the axis of x and component velocities between the above limits respectively, must be proportional to

$$\psi(c) \psi_1(c') du dv dw du' dv' dw' (u - u') dt \dots\dots\dots (A)$$

If the component velocities of these spheres after collision be between

$$U \text{ and } U + dU, \quad V \text{ and } V + dV, \quad W \text{ and } W + dW,$$

$$U' \text{ and } U' + dU', \quad V' \text{ and } V' + dV', \quad W' \text{ and } W' + dW'$$

respectively, then, by the reasoning employed in the last case, we must have

$$\psi(c) \psi_1(c') du \dots dw' (u - u') = \psi(C) \psi_1(C') dU \dots dW' (U' - U), \dots (B)$$

where $V = v, \quad W = w, \quad V' = v', \quad W' = w',$

$$U = u - \frac{2m'}{m+m'}(u - u'),$$

$$U' = u' + \frac{2m}{m+m'}(u - u');$$

whence we get

$$mC^2 + m'C'^2 = mc^2 + m'c'^2,$$

$$\frac{dU}{du'} \frac{dU'}{du} - \frac{dU}{du} \frac{dU'}{du'} = \frac{(m - m')^2}{(m + m')^2} + \frac{4mm'}{(m + m')^2} = 1;$$

and also

$$dV = dv, \quad dW = dw, \quad dV' = dv', \quad dW' = dw',$$

$$U' - U = u - u',$$

$$\text{and} \quad dU dU' = du du' \left(\frac{dU}{du'} \frac{dU'}{du} - \frac{dU}{du} \frac{dU'}{du'} \right) = du du'.$$

So that equation (B) reduces to

$$\psi(c) \psi_1(c') = \psi(C) \psi_1(C'),$$

where

$$mc^2 + m'c'^2 = mC^2 + m'C'^2.$$

A functional equation of which, as before, the solution may be shewn to be

$$\psi(c) = A e^{-\frac{1}{2} mc^2} \quad \text{and} \quad \psi_1(c') = B e^{-\frac{1}{2} m' c'^2}.$$

✓ It must be remembered that h is here a constant of different dimensions from the h of the last proposition.

Since the axis of x may be taken in any direction, it follows that these forms of ψ and ψ_1 will ensure the permanence of distribution as far as collisions between spheres of two different sets are concerned, and we have already proved that they will
 * ensure that permanence so far as collisions between spheres of the same set are concerned; therefore it follows that the number of the N spheres whose component velocities are intermediate between

$$u \text{ and } u+du, \quad v \text{ and } v+dv, \quad w \text{ and } w+dw$$

must be

$$A e^{-h \frac{mc^2}{2}} du dv dw,$$

where

$$c^2 = u^2 + v^2 + w^2;$$

and the number of the N' spheres whose component velocities are intermediate between

$$u' \text{ and } u'+du', \quad v' \text{ and } v'+dv', \quad w' \text{ and } w'+dw'$$

must be

$$B e^{-h \frac{m'c'^2}{2}} du' dv' dw',$$

where

$$c'^2 = u'^2 + v'^2 + w'^2;$$

* ✓ and it is clear that the same reasoning would hold whatever number of sets of such spheres the space might contain.

Applying the results of the last proposition, and writing

$$\frac{mh}{2}, \quad \frac{m'h}{2}$$

instead of h we get

$$A = \frac{N}{\pi^{\frac{3}{2}}} \cdot \left(\frac{mh}{2}\right)^{\frac{3}{2}}, \quad B = \frac{N'}{\pi^{\frac{3}{2}}} \cdot \left(\frac{m'h}{2}\right)^{\frac{3}{2}};$$

and so on for any other sets which the region may contain.

We also find that the mean velocity of each of the N spheres is

$$\frac{2}{\sqrt{\pi}} \cdot \sqrt{\frac{2}{mh}},$$

and similarly for the remaining sets of spheres.

Also the mean square of the velocity of each of the N spheres is

$$\frac{3}{2} \cdot \frac{2}{mh},$$

and the mean vis viva

$$\frac{3}{2h},$$

the last result being the same for each set of spheres.

PROPOSITION III.

On the hypothesis of the last proposition to find the number of pairs of spheres, one being taken from the N set and the other from the N' set, whose relative velocities lie between given limits, and the number of collisions in unit of time and unit of volume between these sets of spheres.

Using the results of the last proposition, writing $\frac{1}{\alpha^2}$ for $\frac{hm}{2}$ and $\frac{1}{\beta^2}$ for $\frac{hm'}{2}$, and supposing the volume of the region considered to be the unit of volume, we find that the number of spheres of the N set which have their component velocities parallel to x between the limits u and $u + du$ is

$$\frac{N}{\alpha\sqrt{\pi}} \cdot e^{-\frac{u^2}{\alpha^2}} du; \dots\dots\dots (A)$$

and that the number of spheres of the N' set which have their component velocities parallel to x between the limits $u + U$ and $u + U + dU$ is

$$\frac{N'}{\beta\sqrt{\pi}} \cdot e^{-\frac{(u+U)^2}{\beta^2}} dU; \dots\dots\dots (B)$$

and therefore the number of pairs of spheres fulfilling the above-mentioned conditions is

$$\frac{NN'}{\alpha\beta\pi} \cdot e^{-\left\{\frac{u^2}{\alpha^2} + \frac{(u+U)^2}{\beta^2}\right\}} du dU.$$

Integrating with respect to u from $-\infty$ to $+\infty$, we find for the total number of pairs of spheres whose relative velocity parallel to x lies between U and $U + dU$ the expression

$$\frac{NN'}{\sqrt{\alpha^2 + \beta^2}} \cdot \frac{1}{\sqrt{\pi}} e^{-\frac{U^2}{\alpha^2 + \beta^2}} dU. \dots\dots\dots (C)$$

Comparing (C) with (A) we see that the distribution of relative velocities follows a law of the same form as that of absolute velocities, and therefore that the mean relative velocity is the square root of the sum of the squares of the mean absolute velocities in the two systems, and the mean square of the relative velocity is the sum of the mean squares of the absolute velocities.

It follows also of course that the number of pairs of spheres, one from each set, whose relative velocities lie between r and $r + dr$ is

$$NN' \frac{4}{\sqrt{\pi} (a^2 + \beta^2)^{\frac{3}{2}}} \cdot e^{-\frac{r^2}{a^2 + \beta^2}} r^2 dr;$$

or, restoring to a and β their values,

$$NN' \frac{\sqrt{2} \cdot (mm'h)^{\frac{3}{2}}}{\sqrt{\pi} (m+m')^{\frac{3}{2}}} \cdot e^{-\frac{r^2 mm'h}{2(m+m')}} r^2 dr.$$

We proceed now to find the number of collisions in unit of time and volume between the spheres of the N set and those of the N' set.

Suppose a number N of equal spheres at rest to be distributed in any manner throughout a unit of volume, and suppose that another sphere moves among them with the velocity r . If a tubular surface be described having for axis the path of the centre of this moving sphere, and for its radius s , or the sum of the radii of the moving sphere and one of the stationary spheres, the volume of the surface thus generated in a unit of time by the moving sphere will be $\pi r s^2$. Hence the chance of the moving sphere colliding with any one of the fixed spheres in a unit of time must be $\pi r s^2$, and the number of collisions in unit of time between the moving sphere and stationary spheres must be $N \pi r s^2$.

The same results would hold good if we replaced the stationary spheres by spheres moving with a common velocity and the moving sphere had a velocity r relative to each of them, that is to say, the chance of collision in unit of time between the last-mentioned sphere and any one of the former-mentioned spheres would be $\pi r s^2$.

Suppose now that there are two sets of N and N' spheres in the unit of volume, and that the number of pairs of spheres (one being taken from each set) whose relative velocities are between r and $r + dr$ is nn' , then, if s be the sum of the radii of each pair, it will follow that the chance of a collision between any pair in unit of time is $\pi r s^2$, and therefore that the total number of collisions in unit of volume and unit of time is $nn' \pi r s^2$.

But we have already seen that nn' is equal to

$$NN' \frac{4}{\sqrt{\pi} (a^2 + \beta^2)^{\frac{3}{2}}} e^{-\frac{r^2}{a^2 + \beta^2}} r^2 dr,$$

N and N' being the total number of spheres of each set in unit-volume, and a^2 and β^2 being written for $\frac{2}{mh}$ and $\frac{2}{m'h}$, as before.

Hence the number of collisions in unit-time between pairs of spheres whose relative velocity lies between r and $r + dr$ is

$$NN' \frac{4\sqrt{\pi}}{(a^2 + \beta^2)^{\frac{3}{2}}} \cdot s^2 r^3 e^{-\frac{r^2}{a^2 + \beta^2}} dr;$$

or
$$NN' \frac{\sqrt{2}}{\sqrt{\pi}} \cdot \frac{(mm'h)^{\frac{3}{2}}}{(m+m')^{\frac{3}{2}}} \cdot s^2 r^3 e^{-\frac{r^2 mm'h}{2(m+m')}} dr.$$

Integrating from $r = 0$ to $r = \infty$, we get

$$2NN' \sqrt{\pi} \cdot \sqrt{a^2 + \beta^2} s^3$$

or
$$2NN' \sqrt{\pi} \sqrt{\frac{2(m+m')}{mm'h}} \cdot s^3$$

for the number of collisions in unit of time which take place in unit of volume between spheres of the N and N' sets.

The number of collisions in the same time and volume between spheres of the N set is therefore

$$2N^2 \sqrt{\pi} \sqrt{2a^2} \cdot s_1^3,$$

or
$$\frac{4N^2 \sqrt{\pi}}{\sqrt{mh}} \cdot s_1^3;$$

and between spheres of the N' set the number is

$$2N'^2 \sqrt{\pi} \sqrt{2\beta^2} s_2^3,$$

or
$$\frac{4N'^2 \sqrt{\pi}}{\sqrt{m'h}} \cdot s_2^3.$$

Also, since the mean velocities in the two systems are

$$\frac{2a}{\sqrt{\pi}} \text{ and } \frac{2\beta}{\sqrt{\pi}},$$

it follows that if l_1 be the mean distance travelled by a sphere of the N_1 set between each collision with a sphere of its own set and l_2 the corresponding mean distance for the second set, then

$$\frac{1}{l_1} = \pi N_1 \sqrt{2} s_1^2 \quad \text{and} \quad \frac{1}{l_2} = \pi N_2 \sqrt{2} s_2^2,$$

s_1 and s_2 being diameters of spheres of the two sets.

Also, if λ_1 be the mean distance travelled by a sphere of the N_1 set without any collision whatever,

$$\frac{1}{\lambda_1} = \pi N_1 \sqrt{2} s_1^2 + \pi N_2 \frac{\sqrt{a^2 + \beta^2}}{a} {}_1s_2^2 + \pi N_3 \frac{\sqrt{a^2 + \gamma^2}}{a} {}_1s_3^2 + \&c.;$$

where ${}_1s_r$ is sum of radii of spheres of the N_1 and N_r sets, with a similar expression for the mean distance travelled by a sphere of any other set.

* * The next case to be considered will be that in which there are any number of sets of spheres as in the last proposition, but acted on by any given impressed forces tending to fixed centres, and therefore capable of being expressed as the space variations of a potential function. Before treating of this case it will be necessary to prove the following general proposition, due originally to Dr. Ludwig Boltzmann and subsequently modified by Professor Maxwell.

Let $p_1 \dots p_n$ be the generalized coordinates, and $q_1 \dots q_n$ the corresponding momenta of any conservative system at any given time, and let $P_1 \dots P_n$, $Q_1 \dots Q_n$ be the corresponding quantities after any interval.

Let the action from the initial to the final state expressed in terms of the initial and final coordinates (p and P) and the energy E be A .

Then, by Thompson and Tait, vol. i, p. 236,

$$q_r = - \frac{dA}{dp_r} \text{ for initial state, } Q_r = \frac{dA}{dP_r} \text{ for final state.}$$

Let $f(p, q)$ be any given function of the p 's and q 's, and let us change the variables in the multiple differential

$$f(p, q) dp_1 \dots dp_n dq_1 \dots dq_n,$$

from the p 's and q 's to the p 's and P 's, i.e. from the $2n$ initial coordinates and momenta to the $2n$ initial and final coordinates.

The expression then becomes

$$f_1(p, P) dp_1 \dots dp_n dP_1 \dots dP_n \Delta;$$

where Δ is the functional determinant

$$\Sigma \pm \left(\frac{dq_1}{dP_1} \dots \frac{dq_n}{dP_n} \right),$$

or

$$\begin{vmatrix} \frac{dq_1}{dP_1} & \dots & \frac{dq_1}{dP_n} \\ \frac{dq_2}{dP_1} & \dots & \frac{dq_2}{dP_n} \\ \vdots & & \vdots \\ \frac{dq_n}{dP_1} & \dots & \frac{dq_n}{dP_n} \end{vmatrix}.$$

Now $\frac{dq_r}{dP_s} = - \frac{d^2 A}{dp_r dP_s} = - \frac{dQ_s}{dp_r};$

$$\therefore \Sigma \pm \left(\frac{dq_1}{dP_1} \dots \frac{dq_n}{dP_n} \right) = - \Sigma \pm \left(\frac{dQ_1}{dp_1} \dots \frac{dQ_n}{dp_n} \right);$$

$$\begin{aligned} \therefore f(p, q) dp_1 \dots dp_n dq_1 \dots dq_n &= f_1(p, P) dp_1 \dots dp_n dP_1 \dots dP_n \Sigma \left(\pm \frac{dq}{dP} \right), \\ &= f_1(p, P) dp_1 \dots dp_n dP_1 \dots dP_n \Sigma \left(\pm \frac{dQ}{dp} \right), \\ &= F(P, Q) dP_1 \dots dP_n dQ_1 \dots dQ_n; \end{aligned}$$

where $F(P, Q)$ and $f(p, q)$ are equal to each other;

$$\therefore dp_1 \dots dp_n dq_1 \dots dq_n = dP_1 \dots dP_n dQ_1 \dots dQ_n;$$

or the product of the differentials remains the same in value after the change of variables.

PROPOSITION IV.

Let there be any number of sets of spheres as in the last proposition, and let them be acted on by any forces passing through the centres of the spheres tending to fixed centres and functions of the distances of the centres of the spheres from the centres of force.

We can no longer assume that the chance of velocities in all directions is the same, as we did in the first two propositions; in other words, we cannot prove, as we did in Proposition I, that the number of spheres of any set having component velocities intermediate between

u and $u+du$, v and $v+dv$, w and $w+dw$ is $\psi(c) du dv dw$,

Let us assume then that the number of spheres of any one set, as the first set, at any time t , having component velocities

between the above-mentioned limits and having the coordinates of their centres intermediate between

$$x \text{ and } x+dx, \quad y \text{ and } y+dy, \quad z \text{ and } z+dz,$$

is

$$\psi(x, y, z, u, v, w) dx dy dz du dv dw.$$

We must in the first place determine the form of

$$\psi(x, y, z, u, v, w),$$

so that it may be independent of t , i.e. so that apart from collisions, the number of spheres of the first set, having their positions and velocities between the limits stated above, may always be the same function of the variables x, y, z, u, v , and w .

✓ Let any sphere whose position and velocity are within the limits defined above be said to be in the state A, and after an interval of time τ let it pass into the state B, such that the coordinates and component velocities are intermediate between

$$X \text{ and } X+dX, \text{ \&c., \&c. } W \text{ and } W+dW.$$

For facility of reference it will be better to denote the states A and B by the subjoined schemes:—

Coordinates of centre between limits.

STATE A. x and $x+dx$, y and $y+dy$, z and $z+dz$;

STATE B. X and $X+dX$, Y and $Y+dY$, Z and $Z+dZ$;

Component velocities between limits.

STATE A. u and $u+du$, v and $v+dv$, w and $w+dw$;

STATE B. U and $U+dU$, V and $V+dV$, W and $W+dW$;

with corresponding changes for the accented variables.

It follows that the number of spheres in the state A at time t must be equal to the number in the state B at time $t+\tau$; and therefore, if the form of ψ be independent of the time,

$$\psi(x, y, z, u, v, w) dx \dots dw = \psi(X, Y, Z, U, V, W) dX \dots dW.$$

✓ But by the last proposition

$$dx \dots dw = dX \dots dW,$$

and therefore

$$\psi(x \dots w) = \psi(X \dots W).$$

Now let

$$\phi_1 = a_1, \quad \phi_2 = a_2, \text{ \&c.}$$

be any equations between the variables $x \dots w$, obtained by the

elimination of t from the integrals of the equations of motion of any one sphere. Then we know that

$$f(\phi_1, \phi_2, \&c.)$$

remains constant throughout the motion.

If therefore $\psi(x \dots w)$ be of the form

$$\psi(\phi_1, \phi_2, \&c.),$$

in other words, if $x, \dots w$ enter into ψ only by entering into $\phi_1, \phi_2, \&c.$, we shall have

$$\psi(x \dots w) = \psi(X \dots W).$$

Hence it follows that, for permanence of distribution independently of collisions, the number of spheres of the N set having the coordinates of their centres and their component velocities intermediate between

$$x \text{ and } x+dx \dots w \text{ and } w+dw$$

must be

$$\psi(\phi_1, \phi_2, \&c.) dx \dots dw.$$

And similarly, the number of spheres of the N' set having the coordinates of their centres and their component velocities intermediate between

$$x' \text{ and } x'+dx' \dots w' \text{ and } w'+dw'$$

must be

$$\psi_1(\phi_1', \phi_2', \&c.) dx' \dots dw'.$$

We must now determine the forms of ψ and ψ_1 in order that this law of distribution may be unaffected by collisions.

By reasoning in all respects the same as in Proposition II we are led to the equation

$$\psi(\phi_1, \phi_2, \&c.) \psi_1(\phi_1', \phi_2', \&c.) = \psi(\Phi_1, \Phi_2, \&c.) \psi_1(\Phi_1', \Phi_2', \&c.),$$

with the equation

$$\frac{mc^2}{2} + \frac{m'c_1'^2}{2} = \frac{mC^2}{2} + \frac{m'C_1'^2}{2}.$$

Also, since the coordinates of the centres are unaffected by collision, it follows that, if χ and χ_1 be the potentials of the impressed forces on the two spheres before collision, and X and X_1 after collision,

$$\chi = X \text{ and } \chi_1 = X_1,$$

$$\text{or } \chi + \frac{mc^2}{2} + \chi_1 + \frac{m'c_1'^2}{2} = X + \frac{mC^2}{2} + X_1 + \frac{m'C_1'^2}{2}.$$

Now the only condition limiting $\phi_1, \phi_2, \&c.$ is that they be equations obtained by eliminating t from any of the integrals

of the equations of motion of the sphere, and the equation of conservation of energy, or

$$\chi + \frac{mc^2}{2} = a,$$

is such an equation, hence ψ may be of the form

$$\psi \left(\chi + \frac{mc^2}{2} \right).$$

Assuming this form for ψ , we get for our condition of permanence

$$\psi \left(\chi + \frac{mc^2}{2} \right) \psi_1 \left(\chi' + \frac{m'c'^2}{2} \right) = \psi \left(X + \frac{mC^2}{2} \right) \psi_1 \left(X' + \frac{m'C'^2}{2} \right),$$

where
$$\chi + \frac{mc^2}{2} + \chi' + \frac{m'c'^2}{2} = X + \frac{mC^2}{2} + X' + \frac{m'C'^2}{2}.$$

A functional equation of which the solution has been proved to be

$$\begin{aligned} \psi &= Ae^{-h \left(\chi + \frac{mc^2}{2} \right)}, \\ \psi_1 &= Be^{-h \left(\chi_1 + \frac{m'c'^2}{2} \right)}; \end{aligned}$$

whence we conclude that the number of spheres of the set N having the coordinates of their centres and their component velocities intermediate between

$$x \text{ and } x+dx \dots w \text{ and } w+dw$$

must be
$$Ae^{-h \left(\chi + \frac{mc^2}{2} \right)} dx dy dz du dv dw,$$

and the number of the set N' similarly circumstanced must be

$$Be^{-h \left(\chi_1 + \frac{m'c'^2}{2} \right)} dx dy dz du dv dw;$$

and so on for all the remaining sets of spheres.

Since the number of spheres of the N set whose centres and velocity points lie within the elementary volumes $dx dy dz$ and $du dv dw$ in the neighbourhood of the points x, y, z , and u, v, w , respectively is

$$Ae^{-h \left(\chi + \frac{mc^2}{2} \right)} du dv dw dx dy dz; \dots \dots \dots (a)$$

where c is the distance of the point u, v, w , from the origin in the velocity diagram, with similar expressions for each of the other sets, it follows that the law of distribution of the velocities and positions of each set of spheres is independent of the co-existence of the other sets in the region under consideration.

If we write, as before, $c^2 dc d\sigma$ for the elementary volume $du dv dw$, the expression (a) becomes

$$Ae^{-h\chi} dx dy dz \cdot e^{-\frac{hmc^2}{2}} c^2 dc d\sigma.$$

Integrating with regard to σ from 0 to 4π and with regard to c from 0 to ∞ , we find that the total number of spheres of any one set whose centres are situated within the elementary volume $dx dy dz$ is given by an expression of the form

$$Be^{-h\chi} dx dy dz. \dots \dots \dots (\beta)$$

That is to say, ρ , or the density of the N set matter in the neighbourhood of the point x, y, z , is

$$\frac{mBe^{-h\chi}}{1}.$$

Again, if we integrate the same expression (a) with regard to x, y , and z for all values corresponding to the region under consideration, we find that the total number of spheres of the N set whose velocity points are situated within the element $dc d\sigma$ is, as in the last case, given by an expression of the form

$$Ce^{-\frac{hmc^2}{2}} c^2 dc d\sigma; \dots \dots \dots (\gamma)$$

whence it follows that so long as the radius vector in the velocity diagram is constant, all directions of that radius vector are equally probable, or that the distribution of velocities is the same in all directions; and this being true of each set of spheres, taken separately, is of course true of the sum total of the spheres in the region under consideration.

Again, integrating (γ) with regard to σ from 0 to 4π , we see that the number of spheres of the N set having velocities between c and $c + dc$ is

$$4\pi Ce^{-\frac{hmc^2}{2}} c^2 dc;$$

whence, as in the last case, we obtain the following results, viz.

The mean velocity of these spheres is $\frac{2\sqrt{2\pi}}{\sqrt{mh}};$

The mean square of the velocities is $\frac{3}{mh};$

✓ The mean vis viva is $\dots \dots \dots \frac{3}{2h}.$

And since the last expression involves only the constant k which is common to all the sets of spheres, it follows that the mean vis viva of the spheres in every set is the same, and therefore equal to the mean vis viva of all the spheres in the region under consideration.

PROPOSITION VI.

A number of sets of elastic spheres are moving about in a given region, as in the last proposition, required to find the action exerted by one portion of the medium on another, or the *pressure*, in the neighbourhood of any point.

Let us assume that the surface of separation of the two portions of the medium is a plane perpendicular to the axis of x , and since the distribution and motion of the spheres of any one set have been proved to be completely independent of the coexistence of the spheres of the remaining sets, let us confine our attention, for the present, to what we have called the N set of spheres.*

Suppose that there are dN such spheres, per unit volume, in the neighbourhood of the point x, y, z , whose component velocities parallel to the axis of x are between u and $u + du$.

The number of these spheres which cross the elementary area $dy dz$ in time dt will be the same as the number of the dN spheres whose centres are situated within the elementary parallelepiped $dx dy dz$ in which dx is equal to $u dt$, and this number is

$$dN \cdot u \cdot dy dz dt.$$

Each of these spheres carries across with it a momentum parallel to the axis of x equal to mu ; the total momentum parallel to the axis of x transferred across $dy dz$ in the time dt is therefore

$$m dN u^2 dy dz dt.$$

If u be positive this is positive momentum transferred from the negative to the positive side of the plane yz , and if u be negative this is negative momentum similarly transferred from the positive to the negative side of the same plane.

* That is to say, the set whereof the number contained in the given region is N .

In either case the result is that by the mere motion of these spheres across the area $dy dz$ the positive momentum parallel to the axis of x is diminished by the quantity $m dN u^2 dy dz dt$ on the negative side of the plane yz and increased by the same quantity on the positive side of that plane in the time dt .

Hence the result is the same as a transference of positive momentum parallel to the axis of x in time dt across the area $dy dz$ equal to $m dy dz dt \sum_{-\infty}^{\infty} u^2 dN$, that is equal to

$$dy dz dt \rho \overline{u^2}; \dots \dots \dots (a)$$

where ρ is the density of the N set matter at the point x, y, z , and $\overline{u^2}$ is the mean square of the velocities parallel to the axis of x .

✓ But either by integration or general reasoning it is easily seen that $\overline{u^2}$ is equal to $\frac{v^2}{3}$, where v^2 is the mean square of the velocities of the N set of spheres, and is equal, as we have seen, to $\frac{3}{mh}$.

Hence there is a transference of positive momentum from the negative to the positive side of the plane yz across the area $dy dz$ in the time dt equal to

$$\rho dy dz dt \frac{v^2}{3}, \text{ or to } \frac{\rho dy dz dt}{mh}.$$

Each separate sphere (whose component velocities are u, v , and w) carries across the same plane momenta parallel to the axes of y and z equal to mv and mw respectively, so that in the time dt there are carried across the elementary area $dy dz$ momenta parallel to the axes of y and z equal to $\sum muv dy dz dt$ and $\sum muw dy dz dt$ respectively.

It is clear from the symmetrical distribution of the velocities that $\sum muv$ and $\sum muw$ are each equal to zero.

Therefore the resultant mutual action of the two portions of the medium across the elementary area $dy dz$ in the time dt is the transference of the momentum $\rho dy dz dt \frac{v^2}{3}$ parallel to the

axis of x across this element from the negative to the positive side.

If this mutual action, or as it is generally called this *pressure*, when referred to unit of surface be denoted by the symbol p , we get the equation

$$p \, dy \, dz \, dt = \rho \, dy \, dz \, dt \cdot \frac{v^2}{3};$$

or

$$p = \rho \frac{v^2}{3} = \frac{\rho}{mh}.$$

Since the momenta parallel to y and z remain unaltered, it follows that the mutual action or pressure between portions of the medium separated by any plane is entirely normal to that plane.

✓ Since also the expression for p , or $\rho \frac{v^2}{3}$, is independent of the direction of the axis of x , it follows that the pressure at any point of the medium is the same in all directions.

If we suppose the contiguous portions of the medium to be separated by a material instead of an ideal plane, it will clearly be necessary for the maintenance of the equilibrium that there should be an action between this plane and the adjacent portion of the medium exactly equivalent to the transference of momenta estimated above. Hence the pressure or force between the plane and medium is normal to the plane, and its value per unit of time and surface is $\frac{\rho}{mh}$ at any point. And this value remains unaffected by turning the plane of separation in any direction about the point.*

* It is important to distinguish between the velocity of agitation of the molecules treated of in the preceding reasoning and that which we are accustomed to consider as the velocity of the medium itself. This latter velocity has been defined by Professor Clerk Maxwell as follows. If we determine the motion of the centre of gravity of all the molecules within a very small region surrounding a point in a medium, then the velocity of the medium within that region is defined as the velocity of the centre of gravity of all the molecules within that region. Should such a velocity exist in the medium under consideration we must suppose that our ideal plane of separation moves with the same velocity, and therefore that the number of molecules crossing it in any direction is on the average equal to the number crossing it in the exactly opposite direction.

We may also prove that the values which we have found for p and ρ are sufficient for the equilibrium of the medium.

For consider the elementary parallelopiped $dx dy dz$ situated in the neighbourhood of the point x, y, z .

The positive momentum of the N sphere matter within this element parallel to the axis of x is, as we have seen, increased by the quantity $p dy dz dt$ in the time dt by transference of matter across the face $dy dz$ nearer to the plane of yz , and the same momentum is, in the same time, diminished by the quantity $(p + \frac{dp}{dx} dx) dy dz dt$ by transference across the opposite face $dy dz$. On the whole, therefore, the positive momentum parallel to x of the N sphere matter within the element is diminished by the quantity $\frac{dp}{dx} dx dy dz dt$ in the time dt .

It is therefore necessary for the permanence of the state of the medium that the impressed force on the N sphere matter within the element should produce a resultant momentum equal to the last found quantity in the time dt , in other words,

$$\frac{dp}{dx} dx dy dz dt \text{ should be equal to } \rho X dx dy dz dt;$$

where X is the impressed force at the point x, y, z parallel to the axis of x ; that is to say,

$$\frac{dp}{dx} \text{ should be equal to } \rho X.$$

$$\text{Now } p = \frac{1}{mh} \rho, \text{ and } \rho = mBe^{-hX};$$

$$\therefore \frac{dp}{dx} = \frac{1}{mh} \frac{d\rho}{dx}, \text{ and } \frac{d\rho}{dx} = -h\rho \frac{dX}{dx} = h\rho mX;$$

$$\therefore \frac{dp}{dx} = \rho X,$$

the required equation.

When several different sets of spheres are present together in the region under consideration the distribution of the centres and of the velocities of the spheres of each set is, as we have seen, independent of the coexistence of the remaining sets. If

therefore $\rho_1, \rho_2, \&c.$ be the densities of the matter of the different sets in the neighbourhood of the point x, y, z , and if $p_1, p_2, \&c.$ be the pressures at that point, defined as above, of the media composed of these different sets, and if $m_1, m_2, \&c.$ be the masses of the individual spheres of each of the sets, and p the total pressure, we shall have

$$\begin{aligned} p &= p_1 + p_2 + \&c. \\ &= \frac{\rho_1}{m_1 h} + \frac{\rho_2}{m_2 h} + \&c. \end{aligned}$$

We will now summarise the results at which we have arrived concerning the physical properties of a medium of sets of elastic spheres moving about in a region of space very much larger than the sum of the volumes of the spheres, and acted on by any impressed forces tending to fixed centres.

In the first place we have found that there is one physical property which has the same value for every set, namely, the mean vis viva of each of the spheres, or $\frac{3}{2h}$. Let us call this quantity τ .

In the second place we have found that the distribution of the positions and velocities of each set of spheres is independent of the remaining sets, and is in all respects the same as if that particular set alone existed in the region of space under consideration.

In the third place we have found that p , the pressure at any point referred to unit of surface, between contiguous portions of the medium arising from the action of any one of the sets is equal to $\frac{2}{3m} \rho \tau$, where ρ is the density of the matter of that particular set in the neighbourhood of the point considered, and m is the mass of each of the spheres of that set, τ being defined as above, and we have also proved that if X, Y , and Z be the component forces at that point, then

$$\frac{dp}{dx} = \rho X, \quad \frac{dp}{dy} = \rho Y, \quad \frac{dp}{dz} = \rho Z.$$

In the fourth place we observe that the fraction $\frac{\rho}{m}$ represents

the number of spheres of the set under consideration per unit of volume in the neighbourhood of the point considered. ✓ It follows

therefore from the equation $p = \frac{2}{3m} \cdot \rho v$ that whenever two media

composed of different sets of spheres are so related that the pressure at any point and the mean vis viva of any one sphere are the same in each medium, then the number of spheres, per unit of volume, is also the same in each medium, or what is the same thing, that the densities of the matter of the two media are

in such a case proportional to the masses of the individual spheres. *

Comparing these results with the well-known properties of a mixture of gases in equilibrium, we know in the first place that there is one physical property, namely the temperature, which is constant throughout such a mixture.

In the second place it has long been recognised as a law established by Dalton that in such a mixture the density of each of the gases at any point is the same as if the other gases

* did not exist, or, as it is frequently expressed, each gas acts as a vacuum to all the rest.

In the third place we know from Boyle's law that so long as the temperature remains the same, the pressure is proportional to the density or inversely proportional to the volume.

In the fourth place we know from the law established by Dalton, Gay Lussac, and Charles, that when the temperature and density both vary, the pressure varies as the temperature and density conjointly.

Finally, we know from Gay Lussac's law that when two gases are at the same pressure and temperature the number of atoms

in unit volume are the same. ✓

We see therefore that in these four particulars the physical properties of a perfect gas, or of a mixture of such gases, exactly correspond with the physical properties of a medium consisting of a set of elastic spheres, or of a mixture of sets of such spheres when the mean vis viva of a sphere is taken as representing the temperature of the gases, or mixture of gases, and the sphere represents the chemical atom. *

Up to this point therefore we have a very strong presump-

tion that the constitution of a homogeneous perfect gas may be explained by supposing it to be a collection of elastic spheres, equal to each other in all respects, and comprised in a region whose volume is very much larger than the sum of the spheres, the temperature of the gas being measured by the mean vis viva of any one of the spheres.

There are however other well-known properties of gases which are not adequately explained on this hypothesis of elastic spheres; the most important of which is the numerical value of the ratio of the specific heats at constant volume and constant pressure respectively.

The quantity of heat required to raise the temperature of unit of mass of any substance one degree is called the specific heat of that substance. In the case of gases this quantity is greater when the gas is allowed to expand under constant pressure than when the volume is kept constant, as when the gas is confined in a rigid vessel, the physical reason of the difference being that in the former case part of the applied heat is converted into mechanical work.

The ratio of the specific heat at constant pressure to that at constant volume has an important bearing on the theory of sound, and has been determined for many gases by careful experiment.

The value of this ratio for several of the more permanent gases, as hydrogen, oxygen, nitrogen, atmospheric air, carbonic oxide, nitrous oxide, and hydrochloric acid, is nearly 1.408; that of the more easily liquifiable gases, as chlorine, ammonia, and sulphuretted hydrogen, is about 1.3; and that of carbonic acid, nitric oxide, sulphurous acid, and marsh gas is about 1.26.

On the hypothesis of smooth elastic spheres this ratio ought to be exactly $1\frac{2}{3}$ or 1.6, as may be seen from the following reasoning:—Suppose r to be the number of such spheres in a unit of mass of the medium, v the volume occupied by such unit, and m the mass of each sphere, then we have the equations

$$mr = 1, \text{ and } pv = 1.$$

Also if τ be the mean vis viva, corresponding to the temperature of the gas, and p the pressure referred to unit of surface,

$$pv = \frac{2}{3}r\tau.$$

Let the mechanical energy δQ in the shape of heat when applied to the medium increase the mean vis viva from τ to $\tau + \delta\tau$, the volume remaining constant, then we must have the equation

$$\delta Q = r\delta\tau.$$

Next let the volume be allowed to vary so that the pressure remains constant, and let $\delta'Q$ be the mechanical energy required to raise the mean vis viva from τ to $\tau + \delta\tau$ in this case.

Since the mass expands under the pressure p , external work equal to $p\delta v$ is performed, δv being the increase of volume, and therefore by the law of conservation of energy

$$\delta'Q = r\delta\tau + p\delta v.$$

But

$$pv = \frac{2}{3}r\tau, \text{ and } p \text{ is constant;}$$

$$\therefore -p\delta v = \frac{2}{3}r\delta\tau;$$

$$\therefore \frac{\delta'Q}{\delta Q} = \frac{1 + \frac{2}{3}}{1} = 1.6.$$

LIBRARY
UNIVERSITY OF
CALIFORNIA.

The only gas for which the ratio has so large a value as this is mercury gas, for which MM. Kundt and Warburg find it to be 1.67.* It is remarkable that the molecule of this substance is regarded by most chemists as monatomic. !!!

There are also other phenomena exhibited by heated gas which suggest a somewhat less simple constitution of the medium than that of moving elastic spheres.

It is known that the light emitted by heated gas, so long as the gas is of no great density, consists of rays of one or more definite kinds of refrangibility, so that when this light is examined by the spectroscope, the spectrum produced consists of one or more bright lines, narrow and distinct, the intervening spaces being dark. As the density of the gas increases, these bright lines become broader and the intervening spaces more luminous, until, as the gas becomes very much condensed, a continuous spectrum is produced. *

If now we replace the moving spheres of the foregoing propositions by a number of bodies all similarly constituted, made up of parts capable of relative motion without being

* See 'Nature' for October, 1875. ✓

altogether torn asunder, this relative motion will be some kind of vibration. The motion of translation or agitation of these small bodies, or molecules, is, as we have seen in the case of the spheres, exceedingly irregular, the intervals between successive encounters and the velocities of a molecule during successive free paths not being subject to any law. It will be different however with the internal motions or vibrations of each molecule. When there is a long free path very many such vibrations may take place in the interval between successive encounters. At each encounter the whole molecule is roughly shaken. During the free path it vibrates according to its own laws, and these vibrations, as is the case in every connected system, may be resolved into a number of simple vibrations, the law of each of which is that of the simple pendulum. At any instant the number of molecules in collision is negligible in comparison with the whole number of molecules in the region under consideration. And therefore at any instant we have a collection of a very great number of bodies, all of which may be regarded as performing precisely the same set of vibrations.

* If these molecules be capable of communicating their vibrations to the luminiferous ether, the result will be light of one or more definite kinds of refrangibility, provided there be any light at all—that is, provided that the vibrations be of such a period as to belong to the luminous part of the spectrum. As the density of this medium is increased, the length of the free path of each molecule is diminished, and since each fresh encounter disturbs the regularity of the series of vibrations, we must no longer regard the whole of the bodies or molecules, but only a large majority of them at any instant, as performing the same sets of vibrations; the result therefore will be light of one or more definite kinds of refrangibility, with a mixture of fainter light of no definite refrangibility, or, viewed under the spectroscope, bright lines of light, along with a ground of diffused light forming a continuous spectrum.*

Let us now proceed to the reconsideration of the problem of the last proposition, the sets of smooth spheres being replaced

* See Maxwell, 'Theory of Heat,' p. 306.

by sets of exactly similar molecules—that is to say, bodies made up of parts which are capable of relative motion without being torn asunder. *

The simplest conception of such bodies would be to regard them as composed of discrete atoms acting upon each other with forces any functions of the distance between them, and in this way the problem has been treated by Dr. Ludwig Boltzmann in his memoirs 'Über das Wärmegleichgewicht zwischen mehratomigen Gasmolekülen,' published in the Transactions of the Imperial Philosophical Academy of Vienna.* For the sake of somewhat greater generality we shall regard our molecules as bodies capable of a given number of degrees of freedom, in other words determined both as to position and arrangement of parts by a given number of generalised coordinates.

These molecules may be acted on by forces either internal, i.e. arising from the mutual action of the different parts of the same system, or external, and directed to fixed centres, but, *in a general way*, each molecule is conceived as being free from the action of any other molecule.

When the distance between any two molecules is so small that they are capable of exerting sensible forces upon one another there will be said to be an encounter between them, such encounters taking the place of the collisions in the case of the spheres, with this difference, that the time of an encounter will not be necessarily indefinitely small. We shall also consider that the forces acting upon the molecules are such that their components may be expressed as the differential coefficients of some function with respect to displacements parallel to the axes of reference, and that they admit therefore of generalised components in terms of the generalised coordinates as explained in Thomson and Tait's 'Natural Philosophy,' vol. i. p. 219. With this explanation we proceed to the consideration of the following proposition. ✓

* Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften Mathematisch-Naturwissenschaftliche Classe, lxiii Band, Zweite Abtheilung, vii Sitzung von 9 März, 1871. *

PROPOSITION VII.

* A number of sets of molecules, the molecules in each set being very numerous and exactly like each other, are in motion in a region of space which is very much larger than the sum of the volumes of all the contained molecules, and are acted on by given forces tending to fixed centres and functions of their distances from these centres, required the law of permanent distribution of coordinates and momenta.

Let the number of generalised coordinates of each of the molecules of one of the given sets be m , and let them be denoted by the symbols $p_1, p_2, \&c., p_m$. Let the corresponding momenta be also denoted by the symbols $q_1, q_2, \&c., q_m$.

Let the total number of molecules of this set be M , and let dM of these be in the state A at any time—that is, let them have their coordinates and momenta intermediate between the values p_1 and $p_1 + dp_1, \&c., p_m$ and $p_m + dp_m, q_1$ and $q_1 + dq_1, \&c., q_m$ and $q_m + dq_m$.

Let also
$$dM = f_m(p_1 \dots p_m, q_1 \dots q_m) dp_1 \dots dq_m.$$

It is required to find the form of f_m in order that it may remain the same for all time.

If $\phi_1 = a_1, \phi_2 = a_2, \&c.$ be any equations obtained by the elimination of t from the integrals of any of the $2m$ equations of motion of one of the M set of molecules, then by reasoning in all respects the same as that employed in Proposition IV it may be proved that so long as there are no encounters the condition of permanent distribution will be satisfied, provided f_m be of the form

$$f_m(\phi_1, \phi_2, \&c.).$$

It remains to find what further restriction must be imposed upon the form of f_m in order that this permanence of arrangement may not be disturbed by any possible encounters.

Suppose that there is another set of N molecules, each of which has n generalised coordinates $r_1, r_2, \&c., r_n$, with the corresponding generalised momenta $s_1, s_2, \&c., s_n$, and let dN of these be in the state B at any time, that is let them have their coordinates and momenta intermediate between r_1 and

$r_1 + dr_1$, &c., r_n and $r_n + dr_n$, s_1 and $s_1 + ds_1$, &c., s_n and $s_n + ds_n$, and suppose that

$$dN = f_n(r_1 \dots r_n s_1 \dots s_n) dr_1 \dots dr_n ds_1 \dots ds_n.$$

Therefore the number of pairs of molecules consisting of one molecule of the set M in state A and another molecule of the set N in state B at any time must be

$$f_m(p_1 \dots p_m q_1 \dots q_m) f_n(r_1 \dots r_n s_1 \dots s_n) dp_1 \dots ds_n.$$

Let $\phi(p_2, \dots, p_m r_1 \dots r_n)$ be such a function of the $m+n-1$ eordinates $p_2 \dots p_m, r_1 \dots r_n$ that whenever p_1 differs from ϕ_1 by less than a certain value a , that is whenever the quantity x in the equation

$$p_1 - \phi = x \dots \dots \dots (1)$$

is equal to or less than a , there is an encounter between the corresponding molecules of the M and N set.

Eliminate p_1 and dp_1 by means of equation (1), and the last found expression becomes

$$f_m(p_1 \dots p_m q_1 \dots q_m) f_n(r_1 \dots r_n s_1 \dots s_n) dx dp_2 \dots dq_m dr_1 \dots ds_n, \dots (2)$$

p_1 being expressed in terms of x and the other variables by means of equation (1). The expression (2) gives the number of pairs of molecules, whereof one is in the state A and the other in the state B at any time: we call such pairs (AB) pairs of molecules.

Let us now introduce the following restriction in the definition of our AB pairs of molecules, namely, that while the values of the $2(m+n)-1$ variables $p_2 \dots s_n$ defining the state of the pair remain as before, the value of the remaining variable p_1 is such as to make x equal to a ; and let us also take dx equal to $\frac{dx}{dt} \delta t$, where $\frac{dx}{dt}$ is formed by the total differentiation of x , or $(p_1 - \phi)$ with respect to t , when the variables in $p_1 - \phi$ have been expressed in terms of t by means of the equations of motion of the molecules. Then it is clear, by reasoning precisely the same as that employed in the elastic spheres, that every one of the (AB) pair of molecules will commence an encounter in the time δt , or in other words the expression

$$f_m \cdot f_n \cdot dp_2 \dots dq_m \cdot dr_1 \dots ds_n \frac{dx}{dt} \cdot \delta t$$

will give us the number of encounters in time δt between a pair

of molecules in the AB state, i. e. with variables intermediate between p_2 and $p_2 + dp_2 \dots s_n$ and $s_n + ds_n$, p_1 being made equal to $\phi + a$ in f_m and $\frac{dx}{dt}$ being found as above explained.

At the end of the time $t + \tau$ let the value of x become again equal to a , i. e. let the encounter terminate, and let the values of $p_2 \dots q_m, r_1 \dots s_n$ at the same instant be $P_2 \dots Q_m, R_1 \dots S_n$.

Then every encounter between a pair of molecules of the M and N set commencing with the variables $p_2 \dots s_n$ being situated within the limits p_2 , and $p_2 + dp_2$, &c., &c., s_n and $s_n + ds_n$, i. e. in the (AB) state, will terminate after an interval τ with the same variables being situated within the limits P_2 and $P_2 + dP_2$, &c., &c., S_n and $S_n + dS_n$, i. e. in the $(A'B')$ state.

Conversely, every encounter commencing between pairs of molecules in the $(A'B')$ state, with the exception only that their momenta have opposite signs to the corresponding momenta in that state, must terminate after the same interval τ in the (AB) state.

It will follow therefore that in order that the permanence of the distribution of the medium may not be affected by an (AB) encounter, the number of such encounters commencing in the time δt must be equal to the number of $(A'B')$ encounters commencing in the same time. But by giving suitable values to the variables $p_2 \dots s_n$, all possible encounters between a molecule of the M set and another of the N set may be represented by an AB encounter, and therefore in order that the permanence of the distribution of the medium may not be affected by any encounter between an M and N molecule the above-mentioned condition must be fulfilled.*

Now we have seen that the number of (AB) encounters commencing in the time δt is equal to

$$f_m(p_1 \dots q_m) f_n(r_1 \dots s_n) dp_2 \dots ds_n \frac{dx}{dt} \cdot \delta t,$$

when we have substituted $\phi + a$ for p_1 and made x equal to a .

✓ * It is assumed, as self-evident, from our supposed constitution of the medium, that the number of pairs of spheres in the $A'B'$ state is the same as the number in that state with the signs of the momenta reversed.

Similarly the number of $A'B'$ encounters is

$$f_m(P_1, \dots, Q_m) f_n(R_1, \dots, S_n) dP_2 \dots dS_n \frac{dX}{dt} \cdot \delta t,$$

when we have substituted $\Phi + X$ for P_1 and made $X = a$.

Hence we must have

$$\begin{aligned} f_m(p_1 \dots q_m) f_n(r_1 \dots s_n) dp_2 \dots ds_n \frac{dx}{dt} \\ = f_m(P_1 \dots Q_m) f_n(R_1 \dots S_n) dP_2 \dots dS_n \frac{dX}{dt}. \end{aligned}$$

Now $dx dp_2 \dots ds_n = dp_1 dp_2 \dots ds_n$.

Also by the general proposition proved above

$$dp_1 dp_2 \dots ds_n = dP_1 \dots dP_2 \dots dS_n;$$

and as before

$$dP_1 dP_2 \dots dS_n = dX dP_2 \dots dS_n;$$

$$\therefore dx dp_2 \dots ds_n = dX dP_2 \dots dS_n.$$

This last equation being true throughout the whole motion it follows that

$$dp_2 \dots ds_n \frac{dx}{dt} = dP_2 \dots dS_n \frac{dX}{dt},$$

and therefore that

$$f_m(p_1 \dots q_m) f_n(r_1 \dots s_n) = f_m(P_1 \dots Q_m) f_n(R_1 \dots S_n).$$

We have seen that f_m and f_n must be of the forms $f_m(\phi_1, \phi_2, \&c.)$, and $f_n(\psi_1, \psi_2, \&c.)$ where $\phi_1 = a_1, \phi_2 = a_2, \&c.$ are equations found by eliminating t from any of the integrals of the $2m$ equations of motion of one of the M set of molecules and $a_1, a_2, \&c.$, are arbitrary constants, while ψ_1, ψ_2 are similarly related to the equations of motion of one of the n set of molecules.

And therefore the equation of condition found above becomes

$$f_m(\phi_1, \phi_2, \&c.) f_n(\psi_1, \psi_2, \&c.) = f_m(\Phi_1, \Phi_2, \&c.) f_n(\Psi_1, \Psi_2, \&c.)$$

Of all the equations of the form $\phi = a$ and $\psi = b$ let us take the equations of conservation of energy and those only, so that f_m and f_n become $f_m(\phi)$ and $f_n(\psi)$ where ϕ and ψ are the sum of the potential and kinetic energies of the respective molecules; then since the total energy, kinetic and potential, in any pair

of molecules must be the same before and after an encounter our equation becomes

$$f_m(\phi)f_n(\psi) = f_m(\Phi)f_n(\Psi), \quad \checkmark$$

subject to the condition

$$\phi + \psi = \Phi + \Psi.$$

Hence, as in the first proposition, we conclude that

$$f_m(\phi) = Ae^{-h\phi} \quad \text{and} \quad f_n(\psi) = Be^{-h\psi};$$

and the expression for dM , or the number of molecules of the M set in the state (A) at any time t , becomes

$$Ae^{-h(\chi+T)} dp_1 \dots dp_m dq_1 \dots dq_m,$$

where χ is the potential and T the kinetic energy corresponding to the state (A) of a molecule. It is obvious that the same law may by the same reasoning be shewn to ensure the permanence of distribution as far as encounters between a pair of molecules from any other pair of sets, or from any similar set, is concerned: it is therefore the law of distribution which we require.

The constant h is the same for every set of molecules, but A may be different for different sets. Hence we see, in the first place, that the law of the arrangement of any one set of molecules is the same as if that set existed alone in the region of space under consideration.

Let the three coordinates p_1, p_2, p_3 be the three rectangular coordinates of the centre of mass of the molecule, and let them be denoted by x, y, z , then if m be the mass of the molecule, and u, v, w be the component velocities of its centre of mass parallel to the same axes, the values of q_1, q_2 , and q_3 will become mu, mv , and mw , so that if $p_4, p_5, \&c. \dots p_m$ be the generalised coordinates of the molecule relative to its centre of mass, and $q_4, q_5, \dots q_m$ the corresponding momenta, the value of T will become

$$\frac{m}{2}(u^2 + v^2 + w^2) + f(p_4 \dots p_m q_4 \dots q_m),$$

✓ where f is the kinetic energy of the molecule relative to its centre of mass, and is therefore an essentially positive quadratic function of the q 's of the form

$$Q_4 q_4^2 + Q_5 q_5^2 + \&c. + R q_4 q_5 + \&c.,$$

the Q 's and R 's being known functions of the p 's.

Therefore the expression for dM , or the number of molecules of the M set in the state A , takes the form of

$$Ce^{-h(x+f)} dx dy dz dp_1 \dots dp_m dq_1 \dots dq_m e^{-h \frac{m}{2}(u^2+v^2+w^2)} du dv dw.$$

If we integrate this with respect to all the independent variables except u , v , and w , we find that the number of molecules of the M set, the component velocities of whose centres of mass parallel to the axes lie between u and $u+du$, v and $v+dv$, w and $w+dw$, is equal to

$$De^{-h \frac{m}{2} \cdot c^2} du dv dw,$$

where $c^2 = u^2 + v^2 + w^2$ and D is independent of u , v , and w .

From this result we infer, precisely as in the case of the elastic spheres,

(1°) That all directions of motion of the centres of mass of the molecules are equally probable.

(2°) That the mean velocity of agitation of a molecule is $\frac{2^{\frac{1}{2}} \cdot \sqrt{\pi}}{\sqrt{mh}}$.

(3°) „ square of the velocity of agitation is $\frac{3}{mh}$.

(4°) „ kinetic energy of agitations parallel to either axis is $\frac{1}{2h}$.

(5°) „ kinetic energy of agitation is $\frac{3}{2h}$.

PROPOSITION VIII.

✓ A number of sets of molecules are in motion in a given region of space, under the circumstances of the last proposition, to find the action of one part of the medium upon another.

The reasoning in this case is in all respects identical with that employed in the corresponding proposition concerning the sets of elastic spheres, and hence we infer

(1) That the pressure or action between two portions of the medium separated by any plane is normal to that plane.

(2) That it is the same for all directions of the normal to that plane.

(3) That the amount of pressure, arising from any one set

of molecules, referred to unit of surface is $\frac{\rho \bar{v}^2}{3}$

where ρ is the density of the matter of which that particular set is composed in the neighbourhood of the point in question, and $\bar{v}^2$ is the mean square of their velocity of translation.

Therefore p , or the pressure referred to unit of surface arising from the action of any one set, is $\frac{\rho}{m} \bar{h}$, and the total value of p is $\Sigma \frac{\rho}{m} \cdot \frac{1}{h}$, where $\Sigma \frac{\rho}{m}$ is the sum of the values of $\frac{\rho}{m}$ for all the sets of molecules.

To find the value of ρ for any one set of molecules in the neighbourhood of any point, we have for the number of molecules of the M set in the state A the expression

$$A e^{-h(\chi+f) + \frac{m}{2}(u^2+v^2+w^2)} dx dy dz du dv dw dp_1 \dots dp_m dq_1 \dots dq_m,$$

where f is a function of the p 's and q 's as already explained.

Integrating with respect to all the momenta the result becomes

$$P. e^{-h(\chi)}. dx dy dz dp_1 \dots dp_m,$$

where P is a function of the p 's only.

Now the forces acting on the molecule are either mutual forces, if any, between different atoms of the molecule or forces tending to fixed centres without the molecule, and we conceive that the dimensions of the molecule are so very small in comparison with the distances of these centres from any parts of it that the latter forces are sensibly parallel and constant. It follows therefore that χ , or the potential energy of the molecule, is the sum of two parts, viz. χ_1 , or the potential energy of the whole mass collected at the centre of mass and acted on by the external forces, and therefore a function of x, y, z only, and χ_2 or the potential energy of the mutual forces (if any), and therefore a function of the p 's only.

Therefore the number of molecules of the M set, whose coordinates lie between x and $x+dx$, &c. ... p_m and p_m+dp_m , is

$$Pe^{-h(\chi_1+\chi_2)}dx dy dz dp_1...dp_m,$$

where P and χ_2 are functions of the p 's only and χ_1 is a function of x, y, z only.

If therefore we integrate the last expression with respect to the p 's we find for the number of molecules of the M set whose centres of mass lie within the parallelopiped $dx dy dz$ the expression

$$Ce^{-h\chi_1}dx dy dz,$$

So that $\rho = m Ce^{-h\chi_1}$ and therefore, as in the elastic sphere case, we obtain the equations

$$\frac{dp}{dx} = \rho X, \quad \frac{dp}{dy} = \rho Y, \quad \frac{dp}{dz} = \rho Z.$$

From the results of this proposition we see, as in the elastic spheres, the agreement between the physical properties of a medium composed of sets of molecules such as we are now considering and a mixture of gases the temperature of which is taken equal to the mean kinetic energy of translation of any one of the molecules. (See pp. 22, 23 above.)

* *

PROPOSITION IX.

A number of sets of molecules are in motion in a given region of space under the circumstances of the two preceding propositions; required to find the mean value of the total kinetic energy of any molecule.

* *

For a molecule of the M set we have seen that the total kinetic energy T is expressed by the equation

$$T = \frac{m}{2}(u^2 + v^2 + w^2) + f,$$

where f is equal to $Q_4q^2_4 + Q_5q^2_5 + \&c. + Rq_4q_5 + \&c.$ as explained above, and we have already proved that the mean value of

the first term in the expression for T is $\frac{3}{2h}$, we proceed now

* to prove that the mean value of f is $\frac{m-3}{2h}$, and therefore that the mean value of T is $\frac{m}{2h}$.

Introduce $m-3$ new variables $r_4, r_5 \dots r_m$ connected with the variables $q_4, q_5 \dots q_m$ by the $m-3$ linear equations

$$\left. \begin{aligned} q_4 &= \lambda_4 r_4 + \lambda_5 r_5 + \&c. + \lambda_m r_m, \\ q_5 &= \mu_4 r_4 + \mu_5 r_5 + \&c. + \mu_m r_m, \\ &\&c., \qquad \&c. \end{aligned} \right\} \dots \dots \dots (a)$$

The λ 's and μ 's being like the Q 's and R 's functions of the $m-3$ coordinates $p_4 \dots p_m$.

Now (Salmon's 'Higher Algebra,' Lesson xii., p. 88) we may determine the λ and μ coefficients in an infinite number of ways, so that the function f may assume the form

$$r_4^2 + r_5^2 + \&c. + r_m^2,$$

the signs preceding each of the terms being positive because no possible values of the q 's, and therefore (by equations a) no possible values of the r 's can make f negative.

Now since dM is the number of molecules of the M set in the state A , at any time it follows that the mean value of such a quantity as r^2 is equal to

$$\frac{\Sigma(r^2 dM)}{\Sigma(dM)}.$$

Eliminate the q 's from the expression for dM found above, i.e.

$$A e^{-h(\chi+T)}. dx \dots dq_m,$$

by means of the equations (a), and it becomes

$$A \Delta e^{-h\chi} dx \dots dp_m \cdot e^{-\frac{mh}{2}(\omega^2 + v^2 + w^2)} du dv dw e^{-h(r_4^2 + \&c. \dots + r_m^2)} dr_4 \dots dr_m,$$

where Δ is the functional determinant

$$\begin{vmatrix} \frac{dq_4}{dr_4} & \dots & \frac{dq_4}{dr_m} \\ \frac{dq_m}{dr_4} & \dots & \frac{dq_m}{dr_m} \end{vmatrix},$$

and therefore independent of the r 's, since the equations (a) are linear. Also, since each of the q 's ranges over all values from $-\infty$ to $+\infty$, it follows from the equations (a) that each of the r 's must admit of a corresponding range.

But $\int_{-\infty}^{\infty} e^{-hr^2} r^2 dr$ is $\frac{\sqrt{\pi}}{2h^{\frac{3}{2}}}$, and $\int_{-\infty}^{\infty} e^{-hr^2} dr$ is $\frac{\sqrt{\pi}}{\sqrt{h}}$.

Therefore

$$\therefore \Sigma r^2 dM = \frac{1}{2h} \left(\frac{\pi}{h}\right)^{\frac{m}{2}} A \iiint \dots \Delta e^{-h\left\{\chi + \frac{m}{2}(u^2+v^2+w^2)\right\}} dx dy dz du dv dw;$$

$$\text{and } \Sigma dM = \left(\frac{\pi}{h}\right)^{\frac{m}{2}} A \iiint \dots \Delta e^{-h\left\{\chi + \frac{m}{2}(u^2+v^2+w^2)\right\}} dx dy dz du dv dw;$$

and therefore the quotient, or the mean value of r^2 , is $\frac{1}{2h}$. *

But the mean value of f is $m-3$ times that of any one of the squares of r . Therefore the mean value of f is $\frac{m-3}{2h}$ and the

mean value of T is $\frac{m}{2h}$. *

PROPOSITION X.

✓ A homogeneous gas being supposed to be constituted of moving molecules with any given number of degrees of freedom, required to find the ratio of the specific heat at constant pressure to that at constant volume. *

First suppose that when a small quantity δQ of heat is imparted to a unit of mass of the gas the temperature is raised from τ to $\tau + \delta\tau$, the volume remaining constant.

Since the volume remains unchanged there is no external work performed, and therefore the whole of this energy δQ must be accounted for by increase of the total kinetic energy and of the potential energy, therefore ✓

$$\delta Q = r(\delta\tau + \delta\chi),$$

where $\delta\tau$ and $\delta\chi$ are the increases of mean total kinetic and

* The result in the text may also be arrived at by direct integration of the expressions

$$\iint \dots A e^{-h(\chi+T)} dx \dots dq_m$$

and

$$\iint \dots A T e^{-h(\chi+T)} dx \dots dq_m$$

over all values of the q 's, where T is an essentially positive homogeneous and quadratic function of these variables.

potential energy respectively of any molecule, and r is the number of molecules in the unit of mass.

We have proved that T , the mean total kinetic energy of a molecule, is equal to $\frac{m\tau}{3}$ where m is the number of degrees of freedom of the molecule;

$$\therefore \delta Q = r \left(\frac{m}{3} + \frac{d\chi}{dt} \right) \delta \tau.$$

✓ Next let the volume be allowed to increase so that the pressure remains constant, and let $\delta'Q$ be the heat required to raise the unit of mass from τ to $\tau + \delta\tau$ in this case.

Since external work is performed equal to $p \delta v$ we must have in this case

$$\delta'Q = r \left(\frac{m}{3} + \frac{d\chi}{dt} \right) \delta \tau + p \delta v.$$

But we know that $p\tau = \frac{2r}{3} \cdot \tau$, and since p remains constant

it follows that $p \delta v = \frac{2r}{3} \delta \tau$;

$$\therefore \frac{\delta'Q}{\delta Q} = \frac{m + 3 \frac{d\chi}{dt} + 2}{m + 3 \frac{d\chi}{dt}}.$$

All that we know of $\frac{d\chi}{dt}$ is that it must necessarily be positive, and of m that it must be integral and not less than 3; writing therefore e for $3 \frac{d\chi}{dt}$ in the calculated value of the ratio of the specific heats, this becomes

$$\frac{m + e + 2}{m + e},$$

or

$$1 + \frac{2}{m + e}.$$

* We have seen that there is one gas for which the ratio is $1\frac{3}{2}$, giving us therefore $m + e = 3$, and thus agreeing with the case of the elastic spheres where there are three degrees of freedom

determined by the coordinates of the centre, and where there is no motion relative to the centre, so that $m = 3$ and $e = 0$.

For the majority of observed gases, however, the ratio is 1.408, giving us

$$\frac{2}{m+e} = 1.408,$$

and

$$m+e = 4.9.$$

For a few gases the ratio is 1.3, whence we get

$$\frac{2}{m+3} = .3,$$

and

$$m+e = 6.6.$$

And for a few other gases the ratio falls as low as 1.26, giving us

$$m+e = 8 \text{ nearly.}$$

The great difficulty in the establishment of the kinetic theory of gases on the molecular hypothesis is to conceive a material system so constituted as to give a suitable value for m in the cases we have mentioned, especially for that most general case of all in which $m+e = 4.9$. ***

Hitherto we have supposed that the gas is so rare, or what is the same thing, that the free path of the molecule is so long, that the number of molecules engaged in encounters at any instant may be neglected in comparison with the number of those which are moving free from such encounters.

On this hypothesis we have found that the number of molecules, each with m generalised coordinates, in any state (A) at any time may be expressed by the formula

$$Ce^{-h(\chi+T)} dp_1 \dots dp_m dq_1 \dots dq_m,$$

where T is the kinetic and χ the potential energy of a molecule whose coordinates and momenta are $p_1, p_2 \dots p_m, q_1, q_2 \dots q_m$ respectively, the forces considered in χ being either the mutual forces between different atoms of the molecule, or external forces tending to fixed centres, but not forces arising from the action of surrounding molecules.

Let us consider now a mass of gas, containing an infinitely

large number of molecules in thermal equilibrium, and suppose that there are r molecules in each unit of mass of this gas.

The chance of any one of these r molecules, as M_1 , being in the state A_1 is as we have seen

$$Ce^{-h(\chi_1 + T_1)} ds_1 d\sigma_1,$$

where ds_1 and $d\sigma_1$ are written for the products $dp_1 \dots dp_m, dq_1 \dots dq_m$ respectively, and χ_1 and T_1 are the values of χ and T when the coordinates and momenta have the values $p_1 \dots p_m, q_1 \dots q_m$ respectively.

Since the total number of molecules in the whole gas is infinitely large, the chance of any one of them being in any particular state may be regarded as independent of the states of the other molecules, and therefore the chance of the r molecules in unit of mass being at any time in the states $A_1, A_2 \dots A_r$ respectively must be

$$Ce^{-h(\chi + T)} ds_1 \dots ds_r, d\sigma_1 \dots d\sigma_r,$$

where $\chi = \chi_1 + \chi_2 \dots + \chi_r$, or the total potential energy of the whole of the r molecules, and $T = T_1 + T_2 \dots T_r$, or the total kinetic energy of the whole of the r molecules in that particular arrangement of the coordinates and velocities.

In this formula, as we have said, intermolecular forces do not enter into the value of χ .

Now it has been proved by Dr. Ludwig Boltzmann in a memoir published in the sixty-fourth volume of the Transactions of the Imperial Philosophical Academy of Vienna, above referred to, that the formula just now obtained will equally
* represent the chance of the above arrangement of the r molecules even when intermolecular forces are taken into account.

In point of fact it is clear that if the r molecules of our unit of mass were to continue free from encounters with any of the remaining molecules, then the above law would ensure a permanent distribution whether intermolecular forces were considered or not, for $\chi + T$ would remain constant, and also the product of the differentials $ds_1 \dots ds_r, d\sigma_1 \dots d\sigma_r$, by what has been proved on p. 12. In fact the reasoning would be in all respects the same as that employed in Prop. IV above, the whole r

molecules together taking the place of any one molecule in that proposition.

✓ In the memoir referred to it is demonstrated that this law of distribution remains unaltered when the effect of encounters between other gas molecules in thermal equilibrium with the given mass of r molecules is taken into consideration.

From this law of distribution it follows, by the same reasoning as has been already employed, that if the mean kinetic energy of agitation of a molecule be taken to represent the temperature τ of the gas, then the mean total kinetic energy of a molecule will be $\frac{m\tau}{3}$ (m being the number of generalised coordinates of the molecule). *

The value of p , or the pressure referred to unit of surface, will however no longer be equal to $\frac{\rho \bar{v}^2}{3}$, as it was in the case where the intermolecular forces were neglected, and before finding its corrected value it will be necessary to prove an important proposition due to Professor Clausius, premising the following definitions. ✓

When any number of material particles are in motion within a limited space so that the particles do not remove continually further and further from their original positions, and provided also that the velocities of the particles do not continually increase or decrease, such a system of material particles is said to be in stationary motion. *

If X, Y, Z be the component forces upon any one of a system of particles in stationary motion, and x, y , and z be the coordinates of such particle referred to any origin and axes, and the quantity $Xx + Yy + Zz$ be found for such particle, then the mean value of the expression

$$-\frac{1}{2} \Sigma (Xx + Yy + Zz)$$

during any period of stationary motion of the system, where Σ represents summation for all the particles, is called the Virial of the system.

PROPOSITION XI.



When a system of material particles is in stationary motion the mean vis viva of the system is equal to its virial.

If x be any function of t ,

$$\frac{d^2}{dt^2}(x^2) = 2 \frac{d}{dt} \cdot (x \frac{dx}{dt}) = 2 (\frac{dx}{dt})^2 + 2x \frac{d^2x}{dt^2};$$

$$\therefore 2 (\frac{dx}{dt})^2 = -2x \frac{d^2x}{dt^2} + \frac{d^2}{dt^2}(x^2). \dots \dots \dots (a)$$

Let x be the coordinate parallel to the axis of x at the time t of that one of the material particles whose mass is m , and let X be the component force on that particle parallel to the same axis, then

$$m \frac{d^2x}{dt^2} = X.$$

Hence equation (a) gives us

$$\frac{m}{2} (\frac{dx}{dt})^2 = -\frac{Xx}{2} + \frac{m}{4} \cdot \frac{d^2}{dt^2}(x^2),$$

$$\therefore \frac{m}{2t} \cdot \int_0^t (\frac{dx}{dt})^2 dt = -\frac{1}{2t} \cdot \int_0^t Xx dt + \frac{m}{4t} \cdot \left[\frac{d(x^2)}{dt} - \left[\frac{d(x^2)}{dt} \right]_0 \right],$$

where $\left[\frac{d(x^2)}{dt} \right]_0$ denotes the initial value of $\frac{d(x^2)}{dt}$.

Now $\frac{1}{t} \cdot \int_0^t (\frac{dx}{dt})^2 dt$ and $\frac{1}{t} \cdot \int_0^t Xx dt$ are the mean values of

$(\frac{dx}{dt})^2$ and Xx during the time t .

✓ If the motion be strictly periodic and t be taken equal to the length of a period or any multiple whatever of that length, then the last term on the right-hand side becomes zero; and if the motion be not strictly periodic, still from its stationary character as above defined, although the coefficient of $\frac{m}{4t}$ in the last term does not necessarily become zero for any value of t however large, yet its value cannot increase indefinitely, but can only fluctuate within certain limits, and therefore by sufficiently

increasing the value of t the last term on the right-hand side becomes inappreciable and may be neglected, so that we obtain, in either case, the equation

$$\frac{m}{2} \cdot \left(\frac{dx}{dt} \right)^2 = -\frac{1}{2} \bar{X}x;$$

and therefore by similar reasoning with respect to y and z , Y and Z ,

$$\frac{m}{2} \cdot \left[\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 + \left(\frac{dz}{dt} \right)^2 \right] = -\frac{1}{2} (\bar{X}x + \bar{Y}y + \bar{Z}z),$$

or
$$\frac{m}{2} \cdot \bar{v}^2 = -\frac{1}{2} (\bar{X}x + \bar{Y}y + \bar{Z}z);$$
 ✓

and therefore for any system of particles,

$$\Sigma \frac{m}{2} \bar{v}^2 = -\frac{1}{2} \cdot \Sigma (\bar{X}x + \bar{Y}y + \bar{Z}z).$$

If the force X , Y , Z arise from the mutual actions of the particles, the expression for the virial may be modified as follows:—

Let R be the force (supposed to be attractive) between any two particles m and m' whose coordinates are xyz , $x'y'z'$, and distance apart r . ✓

Then

$$Xx + X'x' = R \frac{x' - x}{r} \cdot x + R \frac{x - x'}{r} \cdot x' = -R \frac{(x' - x)^2}{r},$$

with similar expressions for $Yy + Y'y'$, $Zz + Z'z'$;

$$\therefore -\frac{1}{2} (Xx + X'x' + Yy + Y'y' + Zz + Z'z') = \frac{1}{2} Rr.$$

Extending this result to any number of particles we find


$$-\frac{1}{2} \Sigma (Xx + Yy + Zz) = \frac{1}{2} \Sigma Rr,$$

and

$$-\frac{1}{2} \Sigma (\bar{X}x + \bar{Y}y + \bar{Z}z) = \frac{1}{2} \Sigma \bar{R}r,*$$

the bar indicating mean values.

* $\Sigma (\bar{R}r)$ represents the mean value of the sum $\Sigma (Rr)$ taken for all possible configurations.



PROPOSITION XII.

✓ A number of material particles are in motion within a closed surface under the action of their mutual forces, required to find the pressure on the surface supposed to be normal to that surface and uniform throughout.

By the last proposition we have

$$\Sigma \frac{m}{2} \overline{v^2} = -\frac{1}{2} \Sigma (\overline{Xx + Yy + Zz}).$$

Also the part of the virial which depends upon mutual forces between the particles has been proved to be equal to

$$\frac{1}{2} \Sigma \overline{Rr},$$

where R is the force (assumed attractive) and r the distance between any two of the particles. It remains to determine that part of the virial which depends upon the normal pressure.

Let p be this pressure referred to unit of surface, and let α, β, γ be the angles which the normal to the surface drawn inwards at any point makes with the axes, and let r be the distance of that point from the origin.

Then the part of the virial arising from the pressure on the element δS of the surface at this point will be

$$-\frac{p}{2} \delta S (x \cos \alpha + y \cos \beta + z \cos \gamma),$$

that is

$$-\frac{p}{2} \cdot r \delta S \cos \theta,$$

where θ is the angle between r and the normal to the surface, drawn inwards.

Conceive a small conical surface to be described, having the origin of coordinates for its vertex, and for its base the element δS of the surface.

If δV be the volume of this elementary cone it is clear that $\frac{\delta S}{2} r \cos \theta$ will be equal in magnitude to $\frac{2}{3} \delta V$, and will be positive or negative according as r meets the surface on the outside or the inside.

Taking the sum of these elementary cones it is clear that the

excess of the negative over the positive sum will be equal to the volume included within the surface.

Hence the whole virial becomes

$$\frac{1}{2}\overline{Rr} + \frac{2}{3}pV,$$

where V is the volume within the surface, and therefore we have this equation for determining p ,

$$pV = \frac{1}{3}\Sigma m\overline{v^2} - \frac{1}{3}\Sigma \overline{Rr}.$$

If the material particles be replaced by molecules it is easy to see that the above-proved property of the virial still holds, when x , y , and z represent the coordinates of the centre of mass of the molecule, and X , Y , and Z are the component forces acting on the molecule supposed to be applied at the centre of mass.

We shall therefore have the equation

$$\Sigma \frac{m}{2}\overline{v^2} = -\frac{1}{2}\Sigma(\overline{Xx} + \overline{Yy} + \overline{Zz}) + \frac{2}{3}pV,$$

where the first term is the part of the virial arising from the mutual forces between the molecules.

It will not be rigorously true for two molecules of any form as it is for two material particles that

$$Xx + X'x' + Yy + Y'y' + Zz + Z'z' = -\frac{1}{2}Rr,$$

R being the mutual force between the molecules (estimated as if the mass of each molecule were collected at its centre of mass), but inasmuch as for the large majority of pairs of molecules their distances apart are infinitely large as compared with their dimensions, and also, in taking the mean value of $Xx + Yy + Zz$, we have to include cases in which the molecules are presented to each other in all manner of relative directions, we conclude, as before, that the mean value of $Xx + Yy + Zz$ may be represented by the mean value of $-Rr$ calculated as if the molecules were material points, so that in all cases we have

$$pV = \frac{2}{3}r\tau - \frac{1}{3}\Sigma Rr,$$

where τ is the temperature, or the mean value of the kinetic energy of translation of a molecule.

When $R=0$, then $pV = \frac{2}{3}r\tau$, agreeing with the result already obtained for the cases in which there are no intermolecular actions.

PROPOSITION XIII.

✓ To deduce the second law of thermodynamics from the results of the kinetic theory of gases.

We have proved that if a gas be composed of an infinite number of molecules, as in the preceding propositions, the chance of any r molecules being in the particular state A at any time may be represented by

$$Ce^{-h(\chi+T)} ds_1 \dots ds_r d\sigma_1 \dots d\sigma_r,$$

where χ and T are the total potential and kinetic energy of the r molecules in the state A , and ds_1 &c. ... $d\sigma_r$, have the meanings assigned to them on p. 40.

Let three of the generalised coordinates in the case of each molecule be the three coordinates of the centre of mass, viz. $x_1 y_1 z_1$ &c., $x_r y_r z_r$, then the above chance may be represented by

$$Ce^{-h(\chi+T)} . dx_1 \dots dz_r . ds . d\sigma,$$

the ds comprising the product of the differentials of all the coordinates except the x , y , and z 's, and the $d\sigma$ comprising the product of the differentials of all the momenta. Also T is of the form $\Sigma \frac{m}{2} (u^2 + v^2 + w^2) + f$, where f is a sum of quadratic functions of the momenta relative to the centres of mass of the different molecules.

Integrating with regard to the momenta, we see that the chance of any configuration may be represented by

$$Pe^{-h\chi} ds . dx_1 \dots dz_r,$$

where P is a function of all the mr coordinates except the $3r$, x , y , and z 's, and ds is the product of the differentials of these $(m-3) \cdot r$ coordinates.

Hence it follows that the mean value $\bar{f}$ of any function f of the positions of the molecules will be determined by the equation

$$\bar{f} = \frac{\iiint \dots f P e^{-h\chi} ds dx_1 \dots dz_r}{\iiint \dots P e^{-h\chi} ds dx_1 \dots dz_r},$$

and that $\bar{\chi}$ or the mean value of the potential energy will be determined by the equation

$$\bar{\chi} = \frac{\iiint \dots \chi P e^{-\lambda \chi} ds dx_1 \dots dz_r}{\iiint \dots P e^{-\lambda \chi} ds dx_1 \dots dz_r}.$$

Suppose now that this unit of mass of gas, consisting of r molecules, is contained in an envelope under the pressure p referred to unit of surface, the volume enclosed by the envelope being v , and let a small accession of heat δQ be imparted to it.

Then h and v will, in general, vary; let their increments be denoted by δh and δv respectively.

There will therefore be an increase of mean kinetic and mean potential energy, the former being $\delta\left(\frac{3r}{2h}\right)$ or $-\frac{3r}{2h^2}\delta h$, and the latter being $\delta\bar{\chi}$, where $\delta\bar{\chi}$ has to be determined by the methods given above.

There will also be external work produced equal to $p\delta v$, and therefore by the conservation of energy we must have

$$\delta Q = -\frac{3r}{2h^2}\delta h + \delta\bar{\chi} + p\delta v;$$

whence we get

$$\frac{\delta Q}{\tau} = -\frac{r}{h}\delta h + \frac{2h}{3}\delta\bar{\chi} + \frac{2h}{3}p\delta v.$$

And the second law of thermodynamics requires that $\frac{\delta Q}{\tau}$, and therefore $-\frac{r}{h}\delta h + \frac{2h}{3}\delta\bar{\chi} + \frac{2h}{3}p\delta v$, should be a perfect differential for all values of δh and δv .

Let us in the first place suppose that there are no intermolecular forces, and let u be taken equal to the expression

$$\log \iiint \dots P e^{-\lambda \chi} ds dx_1 \dots dz_r.$$

Now P and the differentials included in ds' are functions only of the coordinates relative to the centre of mass of the molecules, and are therefore independent of v .

Also since there are no intermolecular forces, it follows that χ is also independent of v .

Therefore
$$\delta u = \frac{du}{dh}\delta h + \frac{du}{dv}\delta v,$$

where the latter term on the right-hand side represents the increase of u arising from the variation of the limits of integration consequent upon the expansion.

Now

$$\frac{du}{dh} \delta h = - \frac{\iint \dots P \chi e^{-h\chi} ds dx_1 \dots dz_r}{\iint \dots P e^{-h\chi} ds dx_1 \dots dz_r} \delta h = -\bar{\chi} \delta h.$$

Also
$$\frac{du}{dv} \delta v = \delta \log \iint \dots (\int P e^{-h\chi} ds) dx_1 \dots dz_r,$$

where δ refers to the variation of the limits of integration in the x, y , and z 's.

Now
$$\begin{aligned} \delta \log \iint \dots (\int P e^{-h\chi} ds) dx_1 \dots dz_r \\ &= \frac{\iint \dots (\int P e^{-h\chi} ds) \delta dx_1 \dots dz_r}{\iint \dots (\int P e^{-h\chi} ds) dx_1 \dots dz_r} \\ &= \frac{\delta \iint \dots (dx_1 \dots dz_r)}{\iint \dots (dx_1 \dots dz_r)} \\ &= \frac{\delta v^r}{v^r} = \frac{r \delta v}{v}; \\ \therefore \quad \frac{r \delta v}{v} &= \bar{\chi} \delta h + \delta u. \end{aligned}$$

Also
$$\frac{\delta Q}{\tau} = -\frac{r}{h} \delta h + \frac{2h}{3} \delta \bar{\chi} + \frac{2}{3} r \delta v.$$

Because $p = \frac{r}{vh}$ in this case

$$\begin{aligned} \therefore \quad \frac{\delta Q}{\tau} &= -\frac{r}{h} \delta v + \frac{2}{3} \delta(h\bar{\chi}) + \frac{2}{3} \delta u \\ &= \delta \left\{ -r \log h + \frac{2}{3} h\bar{\chi} + \frac{2}{3} \log \iint \dots (\int P e^{-h\chi} ds) dx_1 \dots dz_r \right\}. \end{aligned}$$

When the intermolecular forces are considered, the value of $\frac{du}{dv} \delta v$ will consist not only of the term $r \frac{\delta v}{v}$ arising from the variation of the limits of integration with regard to the x, y , and z coordinates, but also of a term arising from the variation

* $\frac{du}{dv} \delta v$ is the alteration which would take place in u , if the volume were increased in the ratio $\frac{v+\delta v}{v}$, the ratio between the coordinates $x_1 \dots z_r$ remaining unaltered.

of χ with regard to v . This latter term is seen from the value of u to be

$$\frac{-h \iint \dots \frac{d\chi}{dv} (\int P e^{-h\chi} ds) dx_1 \dots dz_r}{\iint \dots (\int P e^{-h\chi} ds) dx_1 \dots dz_r} \cdot \delta v;$$

i. e. $-h \frac{d\bar{\chi}}{dv} \delta v;$

and therefore in this case we have

$$\delta u = -\bar{\chi} \delta h + \frac{r \delta v}{v} - h \frac{d\bar{\chi}}{dv} \delta v;$$

$$\therefore \frac{r \delta v}{v} - h \frac{d\bar{\chi}}{dv} \delta v = \bar{\chi} \delta h + \delta u$$

Also in this case we have

$$pv = \frac{r}{h} - \frac{1}{3} \Sigma(\bar{R}r);$$

$$\therefore p \delta v = \frac{r \delta v}{hv} - \frac{\delta v}{3v} \Sigma(\bar{R}r).$$

But if there be a general cubical expansion in the mass from v to $v + \delta v$, this must correspond to a linear expansion from r to $r + \delta r$, where

$$\frac{\delta r}{r} = \frac{1}{3} \cdot \frac{\delta v}{v};$$

$$\therefore p \delta v = \frac{r \delta v}{vh} - \frac{\Sigma R \delta r}{v}.$$

Now since R is the attractive force between any pair of molecules, $\Sigma(R \delta r)$ for any particular configuration must be the increase of potential energy consequent upon the expansion δv in that particular configuration and the mean value of $\Sigma(R \delta r)$ for all possible configurations, or $\overline{\Sigma(R \delta r)}$ must be the mean value of the increases of potential energy for all such configurations, or $\frac{d\bar{\chi}}{dv} \delta v$.*

* It is important to distinguish between $\frac{d\bar{\chi}}{dv}$ or the mean value of the rate of increase of potential energy on expansion for all configurations, and $\frac{d\bar{\chi}}{dv}$ or the rate of increase of the mean potential energy; the former is clearly equal to $\frac{1}{3v} \overline{\Sigma(Rr)}$ by what has been said above, and the latter is found by differentiating the expression for $\bar{\chi}$ with regard to v .

But
$$\frac{\delta Q}{\tau} = -\frac{r}{h} \delta h + \frac{2h}{3} \delta \bar{\chi} + \frac{2h}{3} p \delta v;$$

$$\therefore \frac{\delta Q}{\tau} = -\frac{r}{h} \delta h + \frac{2h}{3} \delta \bar{\chi} + \frac{2r \delta v}{3v} - \frac{2h}{3} \frac{d\bar{\chi}}{dv} \delta v;$$

$$\therefore \frac{\delta Q}{\tau} = -\frac{r}{h} \delta h + \frac{2}{3} (h \delta \bar{\chi} + \bar{\chi} \delta h) + \frac{2}{3} \delta u$$

$$= \delta \left\{ -r \log h + \frac{2}{3} h \bar{\chi} + \frac{2}{3} \log \iint \dots (\int P e^{-h\chi} ds) dx_1 \dots dz_r \right\},$$

as before.

It appears therefore that whether there be appreciable inter-molecular forces, or not, $\frac{\delta Q}{T}$ will be the exact differential of a function ϕ of the independent variables h and v , where ϕ is equal to

$$-r \log h + \frac{2}{3} h \bar{\chi} + \frac{2}{3} \log \iint \dots (\int P e^{-h\chi} ds) dx_1 \dots dz_r.$$

When the molecules become material particles, P must be replaced by unity, and the value of ϕ becomes

$$-r \log h + \frac{2}{3} h \bar{\chi} + \frac{2}{3} \log \iint \dots e^{-h\chi} dx_1 \dots dz_r,$$

this being the form given by Dr. Boltzmann in the memoir referred to above.*

Again, if we write $d\sigma$ for the product $dx_1 \dots dz_r$, we get

$$\bar{\chi} = \frac{\iint \dots \chi P e^{-h\chi} ds d\sigma}{\iint \dots P e^{-h\chi} ds d\sigma};$$

$$\therefore \frac{d\bar{\chi}}{dv} = \frac{\iint \dots \frac{d\chi}{dv} P e^{-h\chi} ds d\sigma}{\iint \dots P e^{-h\chi} ds d\sigma} + \frac{\iint \dots (-h\chi) \frac{d\chi}{dv} P e^{-h\chi} ds d\sigma}{\iint \dots P e^{-h\chi} ds d\sigma}$$

$$+ \frac{\iint \dots \chi P e^{-h\chi} ds d\sigma \iint \dots (-hP \frac{d\chi}{dv}) e^{-h\chi} ds d\sigma}{(\iint \dots P e^{-h\chi} ds d\sigma)^2}.$$

But
$$\frac{d\bar{\chi}}{dv} = \frac{\iint \dots \frac{d\chi}{dv} P e^{-h\chi} ds d\sigma}{\iint \dots P e^{-h\chi} ds d\sigma}.$$

$$\therefore \frac{d}{dh} \left(\frac{d\bar{\chi}}{dv} \right) = \frac{\iint \dots (-\chi) \frac{d\chi}{dv} P e^{-h\chi} ds d\sigma}{\iint \dots P e^{-h\chi} ds d\sigma}$$

$$+ \frac{\iint \dots \chi P e^{-h\chi} ds d\sigma \iint \dots (-P) \frac{d\chi}{dv} e^{-h\chi} ds d\sigma}{(\iint \dots P e^{-h\chi} ds d\sigma)^2};$$

* The proof given above of the second law of thermodynamics on the principles of the kinetic theory of gases is due to Mr. S. H. Burbury, M.A., formerly Fellow of St. John's College, Cambridge, and was published in the Philosophical Magazine for January, 1876.

$$\therefore \frac{d\bar{\chi}}{dv} = \frac{\bar{\chi}}{v} + h \frac{d}{dh} \left(\frac{\bar{\chi}}{dv} \right) = \left(h \frac{d}{dh} + 1 \right) \frac{\bar{\chi}}{dv}.$$

Now if τ be the temperature, we have

$$\tau = \frac{3}{2h};$$

$$\therefore \left(\tau \frac{d}{d\tau} - 1 \right) = - \left(h \frac{d}{dh} + 1 \right).$$

But
$$p = \frac{r}{vh} - \frac{\bar{\chi}}{dv} \text{ (see note p. 49)}$$

$$\therefore - \left(h \frac{d}{dh} + 1 \right) p = - \left(h \frac{d}{dh} + 1 \right) \frac{\bar{\chi}}{dv} = - \frac{d\bar{\chi}}{dv}.$$

But
$$h \frac{d}{dh} + 1 = - \left(\tau \frac{d}{d\tau} - 1 \right);$$

$$\therefore \frac{d\bar{\chi}}{dv} = \tau \frac{dp}{d\tau} - p;$$

$$\therefore \delta \bar{\chi} = \delta \int \left(\tau \frac{dp}{d\tau} - p \right) dv.$$

Now
$$\delta Q = r \delta \tau + \delta \bar{\chi} + p \delta v;$$

therefore
$$\delta Q = r \delta \tau + \delta \cdot \int \left(\tau \frac{dp}{d\tau} - p \right) dv + p \delta v.$$

The expression obtained by Rankine and other writers upon thermodynamics.

May, 1877.

Clarendon Press, Oxford.

BOOKS

PUBLISHED FOR THE UNIVERSITY BY

MACMILLAN AND CO., LONDON;

ALSO TO BE HAD AT THE

CLARENDON PRESS DEPOSITORY, OXFORD.

LEXICONS, GRAMMARS, &c.

(See also Clarendon Press Series pp. 19, 21.)

- A Greek-English Lexicon**, by Henry George Liddell, D.D., and Robert Scott, D.D. *Sixth Edition, Revised and Augmented.* 1870. 4to. cloth, 1l. 16s.
- A Greek-English Lexicon**, abridged from the above, chiefly for the use of Schools. *Seventeenth Edition. Carefully Revised throughout.* 1876. Square 12mo. cloth, 7s. 6d.
- A copious Greek-English Vocabulary**, compiled from the best authorities. 1850. 24mo. bound, 3s.
- Graecae Grammaticae Rudimenta in usum Scholarum.** Auctore Carolo Wordsworth, D.C.L. *Eighteenth Edition,* 1875. 12mo. bound, 4s.
- A Greek Primer**, in English, for the use of beginners. By the Right Rev. Charles Wordsworth, D.C.L., Bishop of St. Andrews. *Fourth Edition.* Extra fcap. 8vo. cloth, 1s. 6d.
- A Practical Introduction to Greek Accentuation**, by H. W. Chandler, M.A. 1862. 8vo. cloth, 10s. 6d.
- Etymologicon Magnum.** Ad Codd. MSS. recensuit et notis variorum instruxit Thomas Gaisford, S.T.P. 1848. fol. cloth, 1l. 12s.
- Suidae Lexicon.** Ad Codd. MSS. recensuit Thomas Gaisford, S.T.P. Tomi III. 1834. fol. cloth, 2l. 2s.

Scheller's Lexicon of the Latin Tongue, with the German explanations translated into English by J. E. Riddle, M.A. 1835. fol. cloth, 1l. 1s.

Scriptores Rei Metricae. Edidit Thomas Gaisford, S.T.P. Tomi III. 8vo. cloth, 15s.

Sold separately:

Hephaestion, Terentianus Maurus, Proclus, cum annotationibus, etc. Tomi II. 10s. Scriptores Latini. 5s.

The Book of Hebrew Roots, by Abu 'L-Walīd Marwān ibn Janāh, otherwise called Rabbi Yōnāh. Now first edited, with an Appendix, by Ad. Neubauer. 4to. cloth, 2l. 7s. 6d.

A Treatise on the use of the Tenses in Hebrew. By S. R. Driver, M.A. Extra fcap. 8vo. cloth, 6s. 6d.

Thesaurus Syriacus : collegerunt Quatremère, Bernstein, Lorsch, Arnoldi, Field: edidit R. Payne Smith, S.T.P.R.
Fasc. I-IV. 1868-77. sm. fol. each, 1l. 1s.

Lexicon Aegyptiaco-Latinum ex veteribus Linguae Aegyptiacae Monumentis, etc., cum Indice Vorum Latinarum ab H. Tattam, A.M. 1835. 8vo. cloth, 15s.

A Practical Grammar of the Sanskrit Language, arranged with reference to the Classical Languages of Europe, for the use of English Students, by Monier Williams, M.A. *Fourth Edition*, 1877. 8vo. cloth, 15s.

A Sanskrit-English Dictionary, Etymologically and Philologically arranged, with special reference to Greek, Latin, German, Anglo-Saxon, English, and other cognate Indo-European Languages. By Monier Williams, M.A., Boden Professor of Sanskrit. 1872. 4to. cloth, 4l. 14s. 6d.

Nalopākhyānam. Story of Nala, an Episode of the Mahā-Bhārata: the Sanskrit text, with a copious Vocabulary, Grammatical Analysis, and Introduction, by Monier Williams, M.A. The Metrical Translation by the Very Rev. H. H. Milman, D.D. 1860. 8vo. cloth, 15s.

Sakuntalā. A Sanskrit Drama, in seven Acts. Edited by Monier Williams, M.A., D.C.L., Boden Professor of Sanskrit. *Second Edition*, 8vo. cloth, 21s.

An Anglo-Saxon Dictionary, by Joseph Bosworth, D.D., Professor of Anglo-Saxon, Oxford. *New edition. In the Press.*

An Icelandic-English Dictionary. Based on the MS. collections of the late Richard Cleasby. Enlarged and completed by G. Vigfússon. With an Introduction, and Life of Richard Cleasby, by G. Webbe Dasent, D.C.L. 4to. cloth, 3l. 7s.

A List of English Words the Etymology of which is illustrated by comparison with Icelandic. Prepared in the form of an APPENDIX to the above. By W. W. Skeat, M.A., *stitched*, 2s.

A Handbook of the Chinese Language. Parts I and II, Grammar and Chrestomathy. By James Summers. 1863. 8vo. *half bound*, 1l. 8s.

Cornish Drama (The Ancient). Edited and translated by E. Norris, Esq., with a Sketch of Cornish Grammar, an Ancient Cornish Vocabulary, etc. 2 vols. 1859. 8vo. *cloth*, 1l. 1s.

The Sketch of Cornish Grammar separately, *stitched*, 2s. 6d.

GREEK CLASSICS, &c.

Aeschylus: quae supersunt in Codice Laurentiano typis descripta. Edidit R. Merkel. 1861. Small folio, *cloth*, 1l. 1s.

Aeschylus: Tragoediae et Fragmenta, ex recensione Guil. Dindorfii. *Second Edition*, 1851. 8vo. *cloth*, 5s. 6d.

Aeschylus: Annotationes Guil. Dindorfii. Partes II. 1841. 8vo. *cloth*, 10s.

Aeschylus: Scholia Graeca, ex Codicibus aucta et emendata a Guil. Dindorfio. 1851. 8vo. *cloth*, 5s.

Sophocles: Tragoediae et Fragmenta, ex recensione et cum commentariis Guil. Dindorfii. *Third Edition*, 2 vols. 1860. fcap. 8vo. *cloth*, 1l. 1s.

Each Play separately, *limp*, 2s. 6d.

The Text alone, printed on writing paper, with large margin, royal 16mo. *cloth*, 8s.

The Text alone, square 16mo. *cloth*, 3s. 6d.

Each Play separately, *limp*, 6d.

Sophocles: Tragoediae et Fragmenta cum Annotatt. Guil. Dindorfii. Tomi II. 1849. 8vo. *cloth*, 10s.

The Text, Vol. I. 5s. 6d. The Notes, Vol. II. 4s. 6d.

Sophocles: Scholia Graeca:

Vol. I. ed. P. Elmsley, A.M. 1825. 8vo. *cloth*, 4s. 6d.

Vol. II. ed. Guil. Dindorfius. 1852. 8vo. *cloth*, 4s. 6d.

Euripides: Tragoediae et Fragmenta, ex recensione Guil. Dindorfii. Tomi II. 1834. 8vo. *cloth*, 10s.

Euripides: Annotationes Guil. Dindorfii. Partes II. 1840. 8vo. *cloth*, 10s.

Euripides: Scholia Graeca, ex Codicibus aucta et emendata a Guil. Dindorfio. Tomi IV. 1863. 8vo. *cloth*, 1l. 16s.

Euripides: *Alcestis*, ex recensione Guil. Dindorfii. 1834. 8vo. *sewed*, 2s. 6d.

- Aristophanes: Comoediae et Fragmenta, ex recensione Guil. Dindorfii.** Tomi II. 1835. 8vo. *cloth*, 11s.
- Aristophanes: Annotationes Guil. Dindorfii.** Partes II. 1837. 8vo. *cloth*, 11s.
- Aristophanes: Scholia Graeca, ex Codicibus aucta et emendata a Guil. Dindorfio.** Partes III. 1839. 8vo. *cloth*, 11.
- Aristophanem, Index in: J. Caravellae.** 1822. 8vo. *cloth*, 3s.
- Metra Aeschyli Sophoclis Euripidis et Aristophanis.** Descripta a Guil. Dindorfio. Accedit Chronologia Scenica. 1842. 8vo. *cloth*, 5s.
- Anecdota Graeca Oxoniensia.** Edidit J. A. Cramer, S.T.P. Tomi IV. 1834-1837. 8vo. *cloth*, 11. 2s.
- Anecdota Graeca e Codd. MSS. Bibliothecae Regiae Parisiensis.** Edidit J. A. Cramer, S.T.P. Tomi IV. 1839-1841. 8vo. *cloth*, 11. 2s.
- Apsinis et Longini Rhetorica.** E Codicibus MSS. recensuit Joh. Bakius. 1849. 8vo. *cloth*, 3s.
- Aristoteles; ex recensione Immanuelis Bekkeri.** Accedunt Indices Sylburgiani. Tomi XI. 1837. 8vo. *cloth*, 21. 10s.
The volumes (except vol. IX.) may be had separately, price 5s. 6d. each.
- Choerobosci Dictata in Theodosii Canones, necnon Epimerismi in Psalmos.** E Codicibus MSS. edidit Thomas Gaisford, S.T.P. Tomi III. 1842. 8vo. *cloth*, 15s.
- Demosthenes: ex recensione Guil. Dindorfii.** Tomi I. II. III. IV. 1846. 8vo. *cloth*, 11. 1s.
- Demosthenes: Tomi V. VI. VII.** Annotationes Interpretum. 1849. 8vo. *cloth*, 15s.
- Demosthenes: Tomi VIII. IX.** Scholia. 1851. 8vo. *cloth*, 10s.
- Harpocrationis Lexicon, ex recensione G. Dindorfii.** Tomi II. 1854. 8vo. *cloth*, 10s. 6d.
- Herculanensium Voluminum Partes II.** 8vo. *cloth*, 10s.
- Homerus: Ilias, cum brevi Annotatione C. G. Heynii.** Accedunt Scholia minora. Tomi II. 1834. 8vo. *cloth*, 15s.
- Homerus: Ilias, ex rec. Guil. Dindorfii.** 1856. 8vo. *cloth*, 5s. 6d.
- Homerus: Scholia Graeca in Iliadem.** Edited by Prof. W. Dindorf, after a new collation of the Venetian MSS. by D. B. Monro, M.A., Fellow of Oriel College. Vols. I. II. 8vo. *cloth*, 24s.
Vols. III. IV. 8vo. *cloth*, 26s. *Just Published.*
- Homerus: Odyssea, ex rec. Guil. Dindorfii.** 8vo. *cloth*, 5s. 6d.

Homerus: Scholia Graeca in Odysseam. Edidit Guil. Dindorfius. Tomi II. 1855. 8vo. *cloth*, 15s. 6d.

Homerum, Index in: Seberi. 1780. 8vo. *cloth*, 6s. 6d.

Oratores Attici ex recensione Bekkeri:

I. Antiphon, Andocides, et Lysias. 1822. 8vo. *cloth*, 7s.

II. Isocrates. 1822. 8vo. *cloth*, 7s.

III. Isaeus, Aeschines, Lycurgus, Dinarchus, etc. 1823. 8vo. *cloth*, 7s.

Scholia Graeca in Aeschinem et Isocratem. Edidit G. Dindorfius. 1852. 8vo. *cloth*, 4s.

Paroemiographi Graeci, quorum pars nunc primum ex Codd. MSS. vulgatur. Edidit T. Gaisford, S.T.P. 1836. 8vo. *cloth*, 5s. 6d.

Plato: The Apology, with a revised Text and English Notes, and a Digest of Platonic Idioms, by James Riddell, M.A. 1867. 8vo. *cloth*, 8s. 6d.

Plato: Philebus, with a revised Text and English Notes, by Edward Poste, M.A. 1860. 8vo. *cloth*, 7s. 6d.

Plato: Sophistes and Politicus, with a revised Text and English Notes, by L. Campbell, M.A. 1866. 8vo. *cloth*, 18s.

Plato: Theaetetus, with a revised Text and English Notes, by L. Campbell, M.A. 1861. 8vo. *cloth*, 9s.

Plato: The Dialogues, translated into English, with Analyses and Introductions, by B. Jowett, M.A., Master of Balliol College and Regius Professor of Greek. A new Edition in 5 volumes, medium 8vo. cloth, 3l. 10s.

Plato: Index to. Compiled for the Second Edition of Professor Jowett's Translation of the Dialogues. By Evelyn Abbott, M.A., Fellow and Tutor of Balliol College. Demy 8vo. *paper covers*, 2s. 6d.

Plato: The Republic, with a revised Text and English Notes, by B. Jowett, M.A., Master of Balliol College and Regius Professor of Greek. Demy 8vo. *Preparing*.

Plotinus. Edidit F. Creuzer. Tomi III. 1835. 4to. 1l. 8s.

Stobaei Florilegium. Ad MSS. fidem emendavit et supplevit T. Gaisford, S.T.P. Tomi IV. 8vo. *cloth*, 1l.

Stobaei Eclogarum Physicarum et Ethicarum libri duo. Accedit Hieroclis Commentarius in aurea carmina Pythagoreorum. Ad MSS. Codd. recensuit T. Gaisford, S.T.P. Tomi II. 8vo. *cloth*, 11s.

Xenophon: Historia Graeca, ex recensione et cum annotationibus L. Dindorfii. *Second Edition*, 1852. 8vo. *cloth*, 10s. 6d.

Xenophon: Expeditio Cyri, ex rec. et cum annotatt. L. Dindorfii. *Second Edition*, 1855. 8vo. *cloth*, 10s. 6d.

Xenophon: Institutio Cyri, ex rec. et cum annotatt. L. Dindorfii. 1857. 8vo. *cloth*, 10s. 6d.

Xenophon: Memorabilia Socratis, ex rec. et cum annotatt. L. Dindorfii. 1862. 8vo. *cloth*, 7s. 6d.

Xenophon: Opuscula Politica Equestria et Venatica cum Arriani Libello de Venatione, ex rec. et cum annotatt. L. Dindorfii. 1866. 8vo. *cloth*, 10s. 6d.

THE HOLY SCRIPTURES, &c.

The Holy Bible in the earliest English Versions, made from the Latin Vulgate by John Wycliffe and his followers: edited by the Rev. J. Forshall and Sir F. Madden. 4 vols. 1850. royal 4to. *cloth*, 3l. 3s.

The Holy Bible: an exact reprint, page for page, of the Authorized Version published in the year 1611. Demy 4to. *half bound*, 1l. 1s.

Vetus Testamentum Graece cum Variis Lectionibus. Editionem a R. Holmes, S.T.P. inchoatam continuavit J. Parsons, S.T.B. Tomi V. 1798–1827. folio, 7l.

Vetus Testamentum ex Versione Septuaginta Interpretum secundum exemplar Vaticanum Romae editum. Accedit potior varietas Codicis Alexandrini. Tomi III. *Editio Altera*. 18mo. *cloth*, 18s.

Origenis Hexaplorum quae supersunt; sive, Veterum Interpretum Graecorum in totum Vetus Testamentum Fragmenta. Edidit Fridericus Field, A.M. 2 vols. 1867–1874. 4to. *cloth*, 5l. 5s.

Libri Psalmorum Versio antiqua Latina, cum Paraphrasi Anglo-Saxonica. Edidit B. Thorpe, F.A.S. 1835. 8vo. *cloth*, 10s. 6d.

Libri Psalmorum Versio antiqua Gallica e Cod. MS. in Bibl. Bodleiana adservato, una cum Versione Metrica aliisque Monumentis pervetustis. Nunc primum descripsit et edidit Franciscus Michel, Phil. Doct. 1860. 8vo. *cloth*, 10s. 6d.

Libri Prophetarum Majorum, cum Lamentationibus Jeremiae, in Dialecto Linguae Aegyptiacae Memphitica seu Coptica. Edidit cum Versione Latina H. Tattam, S.T.P. Tomi II. 1852. 8vo. *cloth*, 17s.

Libri duodecim Prophetarum Minorum in Ling. Aegypt. vulgo Coptica. Edidit H. Tattam, A.M. 1836. 8vo. *cloth*, 8s. 6d.

Novum Testamentum Graece. Antiquissimorum Codicum Textus in ordine parallelo dispositi. Accedit collatio Codicis Sinaitici. Edidit E. H. Hansell, S.T.B. Tomi III. 1864. 8vo. *half morocco*, 2l. 12s. 6d.

Novum Testamentum Graece. Accedunt parallela S. Scripturae loca, necnon vetus capitulorum notatio et canones Eusebii. Edidit Carolus Lloyd, S.T.P.R., necnon Episcopus Oxoniensis. 1873. 18mo. *cloth*, 3s.

The same on writing paper, with large margin, small 4to. *cloth*, 10s. 6d.

Novum Testamentum Graece juxta Exemplar Millianum. 1873. 18mo. *cloth*, 2s. 6d.

The same on writing paper, with large margin, small 4to. *cloth*, 9s.

Evangelia Sacra Graecae. *The Text of Mill.* 1870. fcap. 8vo. *limp*, 1s. 6d.

The New Testament in Greek and English, on opposite pages, arranged and edited by E. Cardwell, D.D. 2 vols. 1837. crown 8vo. *cloth*, 6s.

Novi Testamenti Versio Syriaca Philoxeniana. Edidit Jos. White, S.T.P. Tomi IV. 1278-1803. 4to. *cloth*, 1l. 8s.

Novum Testamentum Coptice, cura D. Wilkins. 1716. 4to. *cloth*, 12s. 6d.

Appendix ad edit. N. T. Gr. e Cod. MS. Alexandrino a C. G. Woide descripti. Subjicitur Codicis Vaticani collatio. 1799. fol. 2l. 2s.

Evangeliorum Versio Gothica, cum Interpr. et Annot. E. Benzeli. Edidit, et Gram. Goth. praemisit, E. Lye, A.M. 1759. 4to. *cloth*, 12s. 6d.

Diatessaron; sive Historia Jesu Christi ex ipsis Evangelistarum verbis apte dispositis confecta. Ed. J. White. 1856. 12mo. *cloth*, 3s. 6d.

Canon Muratorianus. The earliest Catalogue of the Books of the New Testament. Edited with Notes and a Facsimile of the MS. in the Ambrosian Library at Milan, by S. P. Tregelles, LL.D. 1868. 4to. *cloth*, 10s. 6d.

The Five Books of Maccabees, in English, with Notes and Illustrations by Henry Cotton, D.C.L. 1833. 8vo. *cloth*, 10s. 6d.

The Ormulum, now first edited from the original Manuscript in the Bodleian Library (Anglo-Saxon and English), by R. M. White, D.D. 2 vols. *A new Edition in the Press.*

Horae Hebraicae et Talmudicae, a J. Lightfoot. *A new Edition*, by R. Gandell, M.A. 4 vols. 1859. 8vo. *cloth*, 1l. 1s.

FATHERS OF THE CHURCH, &c.

Athanasius: The Orations of St. Athanasius against the Arians. With an Account of his Life. By William Bright, D.D., Regius Professor of Ecclesiastical History, Oxford. Crown 8vo. *cloth*, 9s.

Catenae Graecorum Patrum in Novum Testamentum. Edidit J. A. Cramer, S.T.P. Tomi VIII. 1838-1844. 8vo. *cloth*, 2l. 4s.

- Clementis Alexandrini Opera**, ex recensione Guil. Dindorfii.
Tomi IV. 1869. 8vo. *cloth*, 3*l*.
- Cyrilli Archiepiscopi Alexandrini in XII Prophetas**. Edidit
P. E. Pusey, A.M. Tomi II. 1868. 8vo. *cloth*, 2*l*. 2*s*.
- Cyrilli Archiepiscopi Alexandrini in D. Joannis Evangelium**.
Accedunt Fragmenta Varia necnon Tractatus ad Tiberium Diaconum
Duo. Edidit post Aubertum P. E. Pusey, A.M. Tomi III. 8vo. *cloth*,
2*l*. 5*s*.
- Cyrilli Archiepiscopi Alexandrini Commentarii in Lucae Evan-
gelium** quae supersunt Syriace. E MSS. apud Mus. Britan. edidit R.
Payne Smith, A.M. 1858. 4to. *cloth*, 1*l*. 2*s*.
- The same**, translated by R. Payne Smith, M.A. 2 vols. 1859.
8vo. *cloth*, 1*4s*.
- Ephraemi Syri**, Rabulae Episcopi Edesseni, Balaei, aliorumque.
Opera Selecta. E Codd. Syriacis MSS. in Museo Britannico et Biblio-
theca Bodleiana asservatis primus edidit J. J. Overbeck. 1865. 8vo.
cloth, 1*l*. 1*s*.
- A Latin translation** of the above, by the same Editor. *Pre-
paring*.
- Eusebii Pamphili Evangelicae Praeparationis Libri XV**. Ad
Codd. MSS. recensuit T. Gaisford, S.T.P. Tomi IV. 1843. 8vo.
cloth, 1*l*. 10*s*.
- Eusebii Pamphili Evangelicae Demonstrationis Libri X**. Re-
censuit T. Gaisford, S.T.P. Tomi II. 1852. 8vo. *cloth*, 15*s*.
- Eusebii Pamphili contra Hieroclem et Marcellum Libri**. Re-
censuit T. Gaisford, S.T.P. 1852. 8vo. *cloth*, 7*s*.
- Eusebius' Ecclesiastical History**, according to the text of
Burton. With an Introduction by William Bright, D.D. Crown 8vo.
cloth, 8*s*. 6*d*.
- Eusebii Pamphili Hist. Eccl.: Annotationes Variorum**.
Tomi II. 1842. 8vo. *cloth*, 17*s*.
- Evagrii Historia Ecclesiastica**, ex recensione H. Valesii. 1844.
8vo. *cloth*, 4*s*.
- Heracliti Ephesii Reliquiae**. Recensuit I. Bywater, M.A.
8vo. *cloth*, price 6*s*. *Just Published*.
- Irenaeus: The Third Book of St. Irenaeus**, Bishop of Lyons,
against Heresies. With short Notes, and a Glossary. By H. Deane,
B.D., Fellow of St. John's College, Oxford. Crown 8vo. *cloth*, 5*s*. 6*d*.
- Origenis Philosophumena; sive omnium Haeresium Refutatio**.
E Codice Parisino nunc primum edidit Emmanuel Miller. 1851. 8vo.
cloth, 10*s*.
- Patrum Apostolicorum**, S. Clementis Romani, S. Ignatii, S.
Polycarpi, quae supersunt. Edidit Guil. Jacobson, S.T.P.R. Tomi II.
Fourth Edition, 1863. 8vo. *cloth*, 1*l*. 1*s*.

- Reliquiae Sacrae** secundi tertiiue saeculi. Recensuit M. J. Routh, S.T.P. Tomi V. *Second Edition*, 1846-1848. 8vo. *cloth*, 1l. 5s.
- Scriptorum Ecclesiasticorum Opuscula.** Recensuit M. J. Routh, S.T.P. Tomi II. *Third Edition*, 1858. 8vo. *cloth*, 10s.
- Socratis Scholastici Historia Ecclesiastica.** Gr. et Lat. Edidit R. Hussey, S.T.B. Tomi III. 1853. 8vo. *cloth*, 15s.
- Sozomeni Historia Ecclesiastica.** Edidit R. Hussey, S.T.B. Tomi III. 1859. 8vo. *cloth*, 15s.
- Theodoreti Ecclesiasticae Historiae Libri V.** Recensuit T. Gaisford, S.T.P. 1854. 8vo. *cloth*, 7s. 6d.
- Theodoreti Graecarum Affectionum Curatio.** Ad Codices MSS. recensuit T. Gaisford, S.T.P. 1839. 8vo. *cloth*, 7s. 6d.
- Dowling (J. G.) Notitia Scriptorum SS. Patrum aliorumque vet.** Eccles. Mon. quae in Collectionibus Anecdotorum post annum Christi MDCC. in lucem editis continentur. 1839. 8vo. *cloth*, 4s. 6d.

ECCLESIASTICAL HISTORY, BIOGRAPHY, &c.

- Baedae Historia Ecclesiastica.** Edited, with English Notes, by G. H. Moberly, M.A. 1869. crown 8vo. *cloth*, 10s. 6d.
- Bingham's Antiquities of the Christian Church, and other Works.** 10 vols. 1855. 8vo. *cloth*, 3l. 3s.
- Burnet's History of the Reformation of the Church of England.** *A new Edition.* Carefully revised, and the Records collated with the originals, by N. Pocock, M.A. With a Preface by the Editor. 7 vols. 1865. 8vo. 4l. 4s.
- Burnet's Life of Sir M. Hale, and Fell's Life of Dr. Hammond.** 1856. small 8vo. *cloth*, 2s. 6d.
- Antient Liturgies.** With Introduction, Notes, &c. By C. E. Hammond, M.A. *In Preparation.*
- Cardwell's Two Books of Common Prayer,** set forth by authority in the Reign of King Edward VI, compared with each other. *Third Edition*, 1852. 8vo. *cloth*, 7s.
- Cardwell's Documentary Annals of the Reformed Church of England;** being a Collection of Injunctions, Declarations, Orders, Articles of Inquiry, &c. from 1546 to 1716. 2 vols. 1843. 8vo. *cloth*, 18s.
- Cardwell's History of Conferences on the Book of Common Prayer** from 1551 to 1690. *Third Edition*, 1849. 8vo. *cloth*, 7s. 6d.
- Councils and Ecclesiastical Documents** relating to Great Britain and Ireland. Edited, after Spelman and Wilkins, by A. W. Haddan, B.D., and William Stubbs, M.A., Regius Professor of Modern History, Oxford. Vols. I. and III. Medium 8vo. *cloth*, each 1l. 1s.
- Vol. II. Part I. Medium 8vo. *cloth*, 10s. 6d.

- Formularies of Faith** set forth by the King's Authority during the Reign of Henry VIII. 1856. 8vo. *cloth*, 7s.
- Fuller's Church History** of Britain. Edited by J. S. Brewer, M.A. 6 vols. 1845. 8vo. *cloth*, 1l. 19s.
- Gibson's Synodus Anglicana.** Edited by E. Cardwell, D.D. 1854. 8vo. *cloth*, 6s.
- Hussey's Rise of the Papal Power** traced in three Lectures. *Second Edition*, 1863. fcap. 8vo. *cloth*, 4s. 6d.
- Inett's Origines Anglicanae** (in continuation of Stillingfleet). Edited by J. Griffiths, M.A. 3 vols. 1855. 8vo. *cloth*, 15s.
- John, Bishop of Ephesus.** The Third Part of his Ecclesiastical History. [In Syriac.] Now first edited by William Cureton, M.A. 1853. 4to. *cloth*, 1l. 12s.
- The same**, translated by R. Payne Smith, M.A. 1860. 8vo. *cloth*, 10s.
- Knight's Life of Dean Colet.** 1823. 8vo. *cloth*, 7s. 6d.
- Le Neve's Fasti Ecclesiae Anglicanae.** *Corrected and continued from 1715 to 1853* by T. Duffus Hardy. 3 vols. 1854. 8vo. *cloth*, 1l. 1s.
- Noelli (A.) Catechismus** sive prima institutio disciplinae Pietatis Christianae Latine explicata. Editio nova cura Guil. Jacobson, A.M. 1844. 8vo. *cloth*, 5s. 6d.
- Prideaux's Connection of Sacred and Profane History.** 2 vols. 1851. 8vo. *cloth*, 10s.
- Primers** put forth in the Reign of Henry VIII. 1848. 8vo. *cloth*, 5s.
- Records of the Reformation.** The Divorce, 1527—1533. Mostly now for the first time printed from MSS. in the British Museum and other Libraries. Collected and arranged by N. Pocock, M.A. 2 vols. 8vo. *cloth*, 1l. 16s.
- Reformatio Legum Ecclesiasticarum.** The Reformation of Ecclesiastical Laws, as attempted in the reigns of Henry VIII, Edward VI, and Elizabeth. Edited by E. Cardwell, D.D. 1850. 8vo. *cloth*, 6s. 6d.
- Shirley's (W. W.)** Some Account of the Church in the Apostolic Age. *Second Edition*, 1874. fcap. 8vo. *cloth*, 3s. 6d.
- Shuckford's Sacred and Profane History** connected (in continuation of Prideaux). 2 vols. 1848. 8vo. *cloth*, 10s.
- Stillingfleet's Origines Britannicae**, with Lloyd's Historical Account of Church Government. Edited by T. P. Pantin, M.A. 2 vols. 1842. 8vo. *cloth*, 10s.
- Stubbs's (W.) Registrum Sacrum Anglicanum.** An attempt to exhibit the course of Episcopal Succession in England. 1858. small 4to. *cloth*, 8s. 6d.

Strype's Works Complete, with a General Index. 27 vols.
1821-1843. 8vo. *cloth*, 7l. 13s. 6d. Sold separately as follows:—

Memorials of Cranmer. 2 vols. 1840. 8vo. *cloth*, 11s.
Life of Parker. 3 vols. 1828. 8vo. *cloth*, 16s. 6d.
Life of Grindal. 1821. 8vo. *cloth*, 5s. 6d.
Life of Whitgift. 3 vols. 1822. 8vo. *cloth*, 16s. 6d.
Life of Aylmer. 1820. 8vo. *cloth*, 5s. 6d.
Life of Cheke. 1821. 8vo. *cloth*, 5s. 6d.
Life of Smith. 1820. 8vo. *cloth*, 5s. 6d.
Ecclesiastical Memorials. 6 vols. 1822. 8vo. *cloth*, 1l. 13s.
Annals of the Reformation. 7 vols. 8vo. *cloth*, 2l. 3s. 6d.
General Index. 2 vols. 1828. 8vo. *cloth*, 11s.

Sylloge Confessionum sub tempus Reformandae Ecclesiae editarum. Subjiciuntur Catechismus Heidelbergensis et Canones Synodi Dordrechtanae. 1827. 8vo. *cloth*, 8s.

ENGLISH THEOLOGY.

Beveridge's Discourse upon the XXXIX Articles. *The third complete Edition*, 1847. 8vo. *cloth*, 8s.

Bilson on the Perpetual Government of Christ's Church, with a Biographical Notice by R. Eden, M.A. 1842. 8vo. *cloth*, 4s.

Biscoe's Boyle Lectures on the Acts of the Apostles. 1840. 8vo. *cloth*, 9s. 6d.

Bull's Works, with Nelson's Life. By E. Burton, D.D. *A new Edition*, 1846. 8 vols. 8vo. *cloth*, 2l. 9s.

Burnet's Exposition of the XXXIX Articles. 8vo. *cloth*, 7s.

Burton's (Edward) Testimonies of the Ante-Nicene Fathers to the Divinity of Christ. *Second Edition*, 1829. 8vo. *cloth*, 7s.

Burton's (Edward) Testimonies of the Ante-Nicene Fathers to the Doctrine of the Trinity and of the Divinity of the Holy Ghost. 1831. 8vo. *cloth*, 3s. 6d.

Butler's Works, with an Index to the Analogy. 2 vols. 1874. 8vo. *cloth*, 11s.

Butler's Sermons. 8vo. *cloth*, 5s. 6d.

Butler's Analogy of Religion. 8vo. *cloth*, 5s. 6d.

Chandler's Critical History of the Life of David. 1853. 8vo. *cloth*, 8s. 6d.

Chillingworth's Works. 3 vols. 1838. 8vo. *cloth*, 1l. 1s. 6d.

Clergyman's Instructor. *Sixth Edition*, 1855. 8vo. *cloth*, 6s. 6d.

Comber's Companion to the Temple; or a Help to Devotion in the use of the Common Prayer. 7 vols. 1841. 8vo. *cloth*, 1l. 11s. 6d.

Cranmer's Works. Collected and arranged by H. Jenkyns, M.A., Fellow of Oriel College. 4 vols. 1834. 8vo. *cloth*, 1l. 10s.

Enchiridion Theologicum Anti-Romanum.

Vol. I. Jeremy Taylor's Dissuasive from Popery, and Treatise on the Real Presence. 1852. 8vo. *cloth*, 8s.

Vol. II. Barrow on the Supremacy of the Pope, with his Discourse on the Unity of the Church. 1852. 8vo. *cloth*, 7s. 6d.

Vol. III. Tracts selected from Wake, Patrick, Stillingfleet, Claggett, and others. 1837. 8vo. *cloth*, 11s.

[Fell's] Paraphrase and Annotations on the Epistles of St. Paul. 1852. 8vo. *cloth*, 7s.

Greswell's **Harmonia Evangelica**. *Fifth Edition*, 1856. 8vo. *cloth*, 9s. 6d.

Greswell's **Prolegomena** ad Harmoniam Evangelicam. 1840. 8vo. *cloth*, 9s. 6d.

Greswell's **Dissertations** on the Principles and Arrangement of a Harmony of the Gospels. 5 vols. 1837. 8vo. *cloth*, 3l. 3s.

Hall's (Bp.) **Works**. *A new Edition*, by Philip Wynter, D.D. 10 vols. 1863. 8vo. *cloth*, 3l. 3s.

Hammond's Paraphrase and Annotations on the New Testament. 4 vols. 1845. 8vo. *cloth*, 1l.

Hammond's Paraphrase on the Book of Psalms. 2 vols. 1850. 8vo. *cloth*, 10s.

Heurtley's Collection of Creeds. 1858. 8vo. *cloth*, 6s. 6d.

Homilies appointed to be read in Churches. Edited by J. Griffiths, M.A. 1859. 8vo. *cloth*, 7s. 6d.

Hooker's **Works**, with his Life by Walton, arranged by John Keble, M.A. *Sixth Edition*, 1874. 3 vols. 8vo. *cloth*, 1l. 11s. 6d.

Hooker's **Works**; the text as arranged by John Keble, M.A. 2 vols. 8vo. *cloth*, 11s.

Hooper's (Bp. George) **Works**. 2 vols. 1855. 8vo. *cloth*, 8s.

Jackson's (Dr. Thomas) **Works**. 12 vols. 1844. 8vo. *cloth*, 3l. 6s.

Jewel's **Works**. Edited by R. W. Jelf, D.D. 8 vols. 1847. 8vo. *cloth*, 1l. 10s.

Patrick's **Theological Works**. 9 vols. 1859. 8vo. *cloth*, 1l. 1s.

Pearson's Exposition of the Creed. Revised and corrected by E. Burton, D.D. *Sixth Edition*, 1870. 8vo. *cloth*, 10s. 6d.

Pearson's **Minor Theological Works**. Now first collected, with a Memoir of the Author, Notes, and Index, by Edward Churton, M.A. 2 vols. 1844. 8vo. *cloth*, 10s.

Sanderson's **Works**. Edited by W. Jacobson, D.D. 6 vols. 1854. 8vo. *cloth*, 1l. 10s.

- South's Sermons.** 5 vols. 1842. 8vo. *cloth*, 1l. 10s.
- Stanhope's Paraphrase and Comment upon the Epistles and Gospels.** *A new Edition.* 2 vols. 1851. 8vo. *cloth*, 10s.
- Stillingfleet's Origines Sacrae.** 2 vols. 1837. 8vo. *cloth*, 9s.
- Stillingfleet's Rational Account of the Grounds of Protestant Religion;** being a vindication of Abp. Laud's Relation of a Conference, &c. 2 vols. 1844. 8vo. *cloth*, 10s.
- Wall's History of Infant Baptism, with Gale's Reflections, and Wall's Defence.** *A new Edition*, by Henry Cotton, D.C.L. 2 vols. 1862. 8vo. *cloth*, 1l. 1s.
- Waterland's Works, with Life,** by Bp. Van Mildert. *A new Edition*, with copious Indexes. 6 vols. 1857. 8vo. *cloth*, 2l. 11s.
- Waterland's Review of the Doctrine of the Eucharist, with a Preface** by the present Bishop of London. 1868. crown 8vo. *cloth*, 6s. 6d.
- Wheatly's Illustration of the Book of Common Prayer.** *A new Edition*, 1846. 8vo. *cloth*, 5s.
- Wyclif.** A Catalogue of the Original Works of John Wyclif, by W. W. Shirley, D.D. 1865. 8vo. *cloth*, 3s. 6d.
- Wyclif.** Select English Works. By T. Arnold, M.A. 3 vols. 1871. 8vo. *cloth*, 2l. 2s.
- Wyclif.** Trialogus. *With the Supplement now first edited.* By Gotthardus Lechler. 1869. 8vo. *cloth*, 14s.

ENGLISH HISTORICAL AND DOCUMENTARY WORKS.

- Two of the Saxon Chronicles parallel,** with Supplementary Extracts from the Others. Edited, with Introduction, Notes, and a Glossarial Index, by J. Earle, M.A. 1865. 8vo. *cloth*, 16s.
- Magna Carta,** a careful Reprint. Edited by W. Stubbs, M.A., Regius Professor of Modern History. 1868. 4to. *stitched*, 1s.
- Britton,** a Treatise upon the Common Law of England, composed by order of King Edward I. The French Text carefully revised, with an English Translation, Introduction, and Notes, by F. M. Nichols, M.A. 2 vols. 1865. royal 8vo. *cloth*, 1l. 16s.
- Burnet's History of His Own Time,** with the suppressed Passages and Notes. 6 vols. 1833. 8vo. *cloth*, 2l. 10s.
- Burnet's History of James II,** with additional Notes. 1852. 8vo. *cloth*, 9s. 6d.
- Carte's Life of James Duke of Ormond.** *A new Edition*, carefully compared with the original MSS. 6 vols. 1851. 8vo. *cloth*, 1l. 5s.
- Casauboni Ephemerides,** cum prae fatione et notis J. Russell, S.T.P. Tomi II. 1850. 8vo. *cloth*, 15s.

- Clarendon's** (Edw. Earl of) *History of the Rebellion and Civil Wars in England.* To which are subjoined the Notes of Bishop Warburton. 7 vols. 1849. medium 8vo. *cloth*, 2l. 10s.
- Clarendon's** (Edw. Earl of) *History of the Rebellion and Civil Wars in England.* 7 vols. 1839. 18mo. *cloth*, 1l. 1s.
- Clarendon's** (Edw. Earl of) *History of the Rebellion and Civil Wars in England.* Also *His Life*, written by Himself, in which is included a Continuation of his *History of the Grand Rebellion.* With copious Indexes. In one volume, royal 8vo. 1842. *cloth*, 1l. 2s.
- Clarendon's** (Edw. Earl of) *Life*, including a Continuation of his *History.* 2 vols. 1857. medium 8vo. *cloth*, 1l. 2s.
- Clarendon's** (Edw. Earl of) *Life*, and Continuation of his *History.* 3 vols. 1827. 8vo. *cloth*, 16s. 6d.
- Calendar of the Clarendon State Papers**, preserved in the Bodleian Library.
 Vol. I. From 1523 to January 1649. 8vo. *cloth*, 18s.
 Vol. II. From the death of Charles I, 1649, to the end of the year 1654. 8vo. *cloth*, 16s.
 Vol. III. From 1655 to 1657. 8vo. *cloth*, 14s.
- Freeman's** (E. A.) *History of the Norman Conquest of England: its Causes and Results.* Vols. I, and II. *Second Edition*, with Index. 8vo. *cloth*, 1l. 16s.
 Vol. III. *The Reign of Harold and the Interregnum. Second Edition*, with Index. 8vo. *cloth*, 1l. 1s.
 Vol. IV. *The Reign of William. Second Edition.* 8vo. *cloth*, 1l. 1s.
 Vol. V. *The Effects of the Norman Conquest.* 8vo. *cloth*, 1l. 1s.
- Kennett's** *Parochial Antiquities.* 2 vols. 1818. 4to. *cloth*, 1l.
- Lloyd's** *Prices of Corn in Oxford, 1583-1830.* 8vo. *sewed*, 1s.
- Luttrell's** (Narcissus) *Diary.* A Brief Historical Relation of State Affairs, 1678-1714. 6 vols. 1857. 8vo. *cloth*, 1l. 4s.
- May's** *History of the Long Parliament.* 1854. 8vo. *cloth*, 6s. 6d.
- Rogers's** *History of Agriculture and Prices in England, A.D. 1259-1400.* 2 vols. 1866. 8vo. *cloth*, 2l. 2s.
- Sprigg's** *England's Recovery; being the History of the Army under Sir Thomas Fairfax.* A new edition. 1854. 8vo. *cloth*, 6s.
- Whitelock's** *Memorials of English Affairs from 1625 to 1660.* 4 vols. 1853. 8vo. *cloth*, 1l. 10s.
- Protests of the Lords**, including those which have been expunged, from 1624 to 1874; with Historical Introductions. Edited by James E. Thorold Rogers, M.A. 3 vols. 8vo. *cloth*, 2l. 2s.
- Enactments in Parliament**, specially concerning the Universities of Oxford and Cambridge. Collected and arranged by J. Griffiths, M.A. 1869. 8vo. *cloth* 12s.

Ordinances and Statutes [for Colleges and Halls] framed or approved by the Oxford University Commissioners. 1863. 8vo. *cloth*, 12s.

Sold separately (except for Exeter, All Souls, Brasenose, Corpus, and Magdalen Hall) at 1s. each.

Statuta Universitatis Oxoniensis. 1876. 8vo. *cloth*, 5s.

The Student's Handbook to the University and Colleges of Oxford. *Third Edition.* Extra fcap. 8vo. *cloth*, 2s. 6d.

Index to Wills proved in the Court of the Chancellor of the University of Oxford, &c. Compiled by J. Griffiths, M.A. 1862. royal 8vo. *cloth*, 3s. 6d.

Catalogue of Oxford Graduates from 1659 to 1850. 1851. 8vo. *cloth*, 7s. 6d.

CHRONOLOGY, GEOGRAPHY, &c.

Clinton's Fasti Hellenici. The Civil and Literary Chronology of Greece, from the LVIth to the CXXIIIrd Olympiad. *Third edition*, 1841. 4to. *cloth*, 1l. 14s. 6d.

Clinton's Fasti Hellenici. The Civil and Literary Chronology of Greece, from the CXXIVth Olympiad to the Death of Augustus. *Second edition*, 1851. 4to. *cloth*, 1l. 12s.

Clinton's Epitome of the Fasti Hellenici. 1851. 8vo. *cloth*, 6s. 6d.

Clinton's Fasti Romani. The Civil and Literary Chronology of Rome and Constantinople, from the Death of Augustus to the Death of Heraclius. 2 vols. 1845, 1850. 4to. *cloth*, 3l. 9s.

Clinton's Epitome of the Fasti Romani. 1854. 8vo. *cloth*, 7s.

Cramer's Geographical and Historical Description of Asia Minor. 2 vols. 1832. 8vo. *cloth*, 11s.

Cramer's Map of Asia Minor, 15s.

Cramer's Map of Ancient and Modern Italy, on two sheets, 15s.

Cramer's Description of Ancient Greece. 3 vols. 1828. 8vo. *cloth*, 16s. 6d.

Cramer's Map of Ancient and Modern Greece, on two sheets, 15s.

Greswell's Fasti Temporis Catholici. 4 vols. 1852. 8vo. *cloth*, 2l. 10s.

Greswell's Tables to Fasti, 4to., and Introduction to Tables, 8vo. *cloth*, 15s.

Greswell's Origines Kalendarix Italix. 4 vols. 8vo. *cloth*, 2l. 2s.

Greswell's Origines Kalendarix Hellenicæ. 6 vols. 1862. 8vo. *cloth*, 4l. 4s.

PHILOSOPHICAL WORKS, AND GENERAL LITERATURE.

- The Logic of Hegel**; translated from the *Encyclopaedia of the Philosophical Sciences*. With *Prolegomena*. By William Wallace, M.A. 8vo. *cloth*, 14s.
- Bacon's Novum Organum**, edited, with English notes, by G. W. Kitchin, M.A. 1855. 8vo. *cloth*, 9s. 6d.
- Bacon's Novum Organum**, translated by G. W. Kitchin, M.A. 1855. 8vo. *cloth*, 9s. 6d.
- The Works of George Berkeley, D.D.**, formerly Bishop of Cloyne; including many of his writings hitherto unpublished. With Prefaces, Annotations, and an Account of his Life and Philosophy, by Alexander Campbell Fraser, M.A. 4 vols. 1871. 8vo. *cloth*, 2l. 18s.
- The Life, Letters, &c.** 1 vol. *cloth*, 16s. See also p. 23.
- Smith's Wealth of Nations**. A new Edition, with Notes, by J. E. Thorold Rogers, M.A. 2 vols. 1870. *cloth*, 21s.
- A Course of Lectures on Art**, delivered before the University of Oxford in Hilary Term, 1870. By John Ruskin, M.A., Slade Professor of Fine Art. 8vo. *cloth*, 6s.
- A Critical Account of the Drawings by Michel Angelo and Raffaello** in the University Galleries, Oxford. By J. C. Robinson, F.S.A. Crown 8vo. *cloth*, 4s.

MATHEMATICS, PHYSICAL SCIENCE, &c.

- Archimedis** quae supersunt omnia cum Eutocii commentariis ex recensione Josephi Torelli, cum novâ versione Latinâ. 1792. folio. *cloth*, 1l. 5s.
- Bradley's Miscellaneous Works and Correspondence**. With an Account of Harriot's Astronomical Papers. 1832. 4to. *cloth*, 17s.
Reduction of Bradley's Observations by Dr. Busch. 1838. 4to. *cloth*, 3s.
- Treatise on Infinitesimal Calculus**. By Bartholomew Price, M.A., F.R.S., Professor of Natural Philosophy, Oxford.
- Vol. I. Differential Calculus. *Second Edition*, 1858. 8vo. *cloth*, 14s. 6d.
 - Vol. II. Integral Calculus, Calculus of Variations, and Differential Equations. *Second Edition*, 1865. 8vo. *cloth*, 18s.
 - Vol. III. Statics, including Attractions; Dynamics of a Material Particle. *Second Edition*, 1868. 8vo. *cloth*, 16s.
 - Vol. IV. Dynamics of Material Systems; together with a Chapter on Theoretical Dynamics, by W. F. Donkin, M.A., F.R.S. 1862. 8vo. *cloth*, 16s.

- A Treatise on the Kinetic Theory of Gases.** By Henry William Watson, M.A., formerly Fellow of Trinity College, Cambridge. 1876. 8vo. *cloth*, 3s. 6d.
- Rigaud's Correspondence of Scientific Men of the 17th Century,** with Index by A. de Morgan. 2 vols. 1841-1862. 8vo. *cloth*, 18s. 6d.
- Daubeny's Introduction to the Atomic Theory.** *Second Edition*, greatly enlarged. 1850. 16mo. *cloth*, 6s.
- Vesuvius.** By John Phillips, M.A., F.R.S., Professor of Geology, Oxford. 1869. Crown 8vo. *cloth*, 10s. 6d.
- Geology of Oxford and the Valley of the Thames.** By the same Author. 8vo. *cloth*, 21s.
- Synopsis of the Pathological Series in the Oxford Museum.** By H. W. Acland, M.D., F.R.S., 1867. 8vo. *cloth*, 2s. 6d.
- Thesaurus Entomologicus Hopeianus, or a Description of the rarest Insects in the Collection given to the University by the Rev. William Hope.** By J. O. Westwood, M.A. With 40 Plates, mostly coloured. Small folio, *half morocco*, 7l. 10s.
- Text-Book of Botany, Morphological and Physiological.** By Dr. Julius Sachs, Professor of Botany in the University of Würzburg. Translated by A. W. Bennett, M.A. Royal 8vo. *half morocco*, 1l. 11s. 6d.

BIBLIOGRAPHY.

- Ebert's Bibliographical Dictionary**, translated from the German. 4 vols. 1837. 8vo. *cloth*, 1l. 10s.
- Cotton's List of Editions of the Bible in English.** *Second Edition*, corrected and enlarged. 1852. 8vo. *cloth*, 8s. 6d.
- Cotton's Typographical Gazetteer.** 1831. 8vo. *cloth*, 12s. 6d.
- Cotton's Typographical Gazetteer, Second Series.** 1866. 8vo. *cloth*, 12s. 6d.
- Cotton's Rhemes and Doway.** An attempt to shew what has been done by Roman Catholics for the diffusion of the Holy Scriptures in English. 1855. 8vo. *cloth*, 9s.

BODLEIAN LIBRARY CATALOGUES, &c.

Catalogus Codd. MSS. Orientalium Bibliothecae Bodleianae :

Pars I, a J. Uri. 1788. fol. 10s.

Partis II Vol. I, ab A. Nicoll, A.M. 1821. fol. 10s.

Partis II Vol. II, Arabicos complectens, ab E. B. Pusey, S.T.B. 1835. fol. 1l.

Catalogus MSS. qui ab E. D. Clarke comparati in Bibl. Bodl. adservantur :

Pars prior. Inseruntur Scholia inedita in Platonem et in Carmina Gregorii Naz. 1812. 4to. 5s.

Pars posterior, Orientales complectens, ab A. Nicoll, A.M. 1814. 4to. 2s. 6d.

Catalogus Codd. MSS. et Impressorum cum notis MSS. olim D'Orvillianorum, qui in Bibl. Bodl. adservantur. 1806. 4to. 2s. 6d.

Catalogus MSS. Borealium praecipue Islandicae Originis, a Finno Magno Islando. 1832. 4to. 4s.

Catalogus Codd. MSS. Bibliothecae Bodleianae:—

Pars I. Codices Graeci, ab H. O. Coxe, A.M. 1853. 4to. 1l.

Partis II. Fasc. I. Codices Laudiani, ab H. O. Coxe, A.M. 1858. 4to. 1l.

Pars III. Codices Graeci et Latini Canonici, ab H. O. Coxe, A.M. 1854. 4to. 1l.

Pars IV. Codices T. Tanneri, ab A. Hackman, A.M. 1860. 4to. 12s.

Pars V. Codicum R. Rawlinson classes duae priores, a Guil. D. Macray, A.M. 1862. 4to. 12s.

Pars VI. Codices Syriaci, a R. P. Smith, A.M. 1864. 4to. 1l.

Pars VII. Codices Aethiopici, ab A. Dillmann, Ph. Doct. 1848. 4to. 6s. 6d.

Pars VIII. Codices Sanscritici, a Th. Aufrecht, A.M. 1859-1864. 4to. 1l. 10s.

Catalogo di Codici MSS. Canonici Italiani, compilato dal Conte A. Mortara. 1864. 4to. 10s. 6d.

Catalogus Librorum Impressorum Bibliothecae Bodleianae. Tomi IV. 1843 to 1850. fol. 4l.

Catalogus Dissertationum Academicarum quibus nuper aucta est Bibliotheca Bodleiana. 1834. fol. 7s.

Catalogue of Books bequeathed to the Bodleian Library by R. Gough, Esq. 1814. 4to. 15s.

Catalogue of Early English Poetry and other Works illustrating the British Drama, collected by Edmond Malone, Esq. 1835. fol. 4s.

Catalogue of the Printed Books and Manuscripts bequeathed to the Bodleian Library by Francis Douce, Esq. 1840. fol. 15s.

Catalogue of a Collection of Early Newspapers and Essayists presented to the Bodleian Library by the late Rev. F. W. Hope. 1865. 8vo. 7s. 6d.

Catalogue of the Manuscripts bequeathed to the University of Oxford by Elias Ashmole. By W. H. Black. 1845. 4to. 1l. 10s.

Index to the above, by W. D. Macray, M.A. 1867. 4to. 10s.

Catalogus Codd. MSS. qui in Collegiis Aulisque Oxoniensibus hodie adservantur. Confecit H. O. Coxe, A.M. Tomi II. 1852. 4to. 2l.

Catalogus Codd. MSS. in Bibl. Aed. Christi ap. Oxon. Curavit G. W. Kitchin, A.M. 1867. 4to. 6s. 6d.

Clarendon Press Series.

The Delegates of the Clarendon Press having undertaken the publication of a series of works, chiefly educational, and entitled the *Clarendon Press Series*, have published, or have in preparation, the following.

Those to which prices are attached are already published; the others are in preparation.

I. LATIN; GRAMMAR, CLASSICS, &c.

An Elementary Latin Grammar. By John B. Allen, M.A., Head Master of Perse Grammar School, Cambridge. Extra fcap. 8vo. cloth, 2s. 6d.

A First Latin Exercise Book. By the same Author. Extra fcap. 8vo. cloth, 2s. 6d.

A Series of Graduated Latin Readers.

First Latin Reader. By T. J. Nunns, M.A. *Second Edition.* Extra fcap. 8vo. cloth, 2s.

Second Latin Reader.

Third Latin Reader, or Specimens of Latin Literature. Part I, Poetry. By James McCall Marshall, M.A., Dulwich College.

Fourth Latin Reader.

Cicero. Selection of interesting and descriptive passages. With Notes. By Henry Walford, M.A., Wadham College, Oxford, Assistant Master at Haileybury College. In three Parts. *Second Edition.* Extra fcap. 8vo. cloth, 4s. 6d.

Each Part separately, limp, 1s. 6d.

Part I. Anecdotes from Grecian and Roman History.

Part II. Omens and Dreams: Beauties of Nature.

Part III. Rome's Rule of her Provinces.

Cicero. Selected Letters (for Schools). With Notes. By the late C. E. Prichard, M.A., formerly Fellow of Balliol College, Oxford, and E. R. Bernard, M.A., Fellow of Magdalen College, Oxford. *Second Edition.* Extra fcap. 8vo. cloth, 3s.

Pliny. Selected Letters (for Schools). With Notes. By the same Editors. Extra fcap. 8vo. cloth, 3s.

Caesar. The Commentaries (for Schools). With Notes and Maps. By Charles E. Moberly, M.A.

Part I. The Gallic War. *Third Edition.* Extra fcap. 8vo. *cloth*, 4s. 6d.

Part II. The Civil War. Extra fcap. 8vo. *cloth*, 3s. 6d.

The Civil War. Book I. Extra fcap. 8vo. *cloth*, 2s.

Cornelius Nepos. With Notes. By Oscar Browning, M.A., Fellow of King's College, Cambridge, and Assistant Master at Eton College. Extra fcap. 8vo. *cloth*, 2s. 6d.

Livy. Selections (for Schools). With Notes and Maps. By H. Lee-Warner, M.A., Assistant Master in Rugby School. Extra fcap. 8vo. *In Parts, limp*, each 1s. 6d.

Part I. The Caudine Disaster.

Part II. Hannibal's Campaign in Italy.

Part III. The Macedonian War.

Livy, Books I-X. By J. R. Seeley, M.A., Regius Professor of Modern History, Cambridge. Book I. *Second Edition.* 8vo. *cloth*, 6s.

Also a small edition for Schools.

Tacitus. The Annals. Books I-VI. With Essays and Notes. By T. F. Dallin, M.A., Tutor of Queen's College, Oxford. *Preparing.*

Passages for Translation into Latin. For the use of Passmen and others. Selected by J. Y. Sargent, M.A., Fellow and Tutor of Magdalen College, Oxford. *Third Edition.* Ext. fcap. 8vo. *cloth*, 2s. 6d.

Cicero's Philippic Orations. With Notes. By J. R. King, M.A., formerly Fellow and Tutor of Merton College. 8vo. *cloth*, 10s. 6d.

Cicero. Select Letters. With English Introductions, Notes, and Appendices. By Albert Watson, M.A., Fellow and formerly Tutor of Brasenose College, Oxford. *Second Edition.* Demy 8vo. *cloth*, 18s.

Cicero. Select Letters. Text. By the same Editor. Extra fcap. 8vo. *cloth*, 4s.

Cicero pro Cluentio. With Introduction and Notes. By W. Ramsay, M.A. Edited by G. G. Ramsay, M.A., Professor of Humanity, Glasgow. Extra fcap. 8vo. *cloth*, 3s. 6d.

Cicero de Oratore. With Introduction and Notes. By A. S. Wilkins, M.A., Professor of Latin, Owens College, Manchester.

Catulli Veronensis Liber. Recognovit, apparatus criticum prolegomena appendices addidit, Robinson Ellis, A.M. 1867. 8vo. *cloth*, 16s.

Catulli Veronensis Carmina Selecta, secundum recognitionem Robinson Ellis, A.M. Extra fcap. 8vo. *cloth*, 3s. 6d.

A Commentary on Catullus. By Robinson Ellis, M.A. Demy 8vo. *cloth*, 16s.

Horace. With a Commentary. Volume I. The Odes, Carmen Seculare, and Epodes. By Edward C. Wickham, M.A., Head Master of Wellington College. 8vo. cloth, 12s.

Also a small edition for Schools.

Ovid. Selections for the use of Schools. With Introductions and Notes, and an Appendix on the Roman Calendar. By W. Ramsay, M.A. Edited by G. G. Ramsay, M.A., Professor of Humanity, Glasgow. *Second Edition.* Ext. fcap. 8vo. cloth, 5s. 6d.

Persius. The Satires. With a Translation and Commentary. By John Conington, M.A., late Corpus Professor of Latin in the University of Oxford. Edited by Henry Nettleship, M.A. *Second Edition.* 8vo. cloth, 7s. 6d.

Vergil: Suggestions Introductory to a Study of the Aeneid. By H. Nettleship, M.A. 8vo. sewed, 1s. 6d.

Selections from the less known Latin Poets. By North Pinder, M.A. Demy 8vo. cloth, 15s.

Fragments and Specimens of Early Latin. With Introductions and Notes. By John Wordsworth, M.A., Tutor of Brasenose College, Oxford. Demy 8vo. cloth, 18s.

A Manual of Comparative Philology. By T. L. Papillon, M.A., Fellow and Lecturer of New College. Crown 8vo. cloth, 6s.

The Ancient Languages of Italy. By Theodore Aufrecht, Phil. Doct. *Preparing.*

The Roman Poets of the Augustan Age. By William Young Sellar, M.A., Professor of Humanity in the University of Edinburgh. VIRGIL. 8vo. cloth, 14s.

The Roman Poets of the Republic. By the same Editor. *Preparing.*

II. GREEK; GRAMMARS, CLASSICS, &c.

A Greek Primer in English for the use of beginners. By the Right Rev. Charles Wordsworth, D.C.L., Bishop of St. Andrews. *Fourth Edition.* Extra fcap. 8vo. cloth, 1s. 6d.

Graecae Grammaticae Rudimenta in usum Scholarum. Auctore Carolo Wordsworth, D.C.L. *Eighteenth Edition,* 1875. 12mo. bound, 4s.

A Greek-English Lexicon, abridged from Liddell and Scott's 4to. edition, chiefly for the use of Schools. *Seventeenth Edition.* *Carefully Revised throughout.* 1876. Square 12mo. cloth, 7s. 6d.

Greek Verbs, Irregular and Defective; their forms, meaning, and quantity; embracing all the Tenses used by Greek writers, with reference to the passages in which they are found. By W. Veitch. *New Edition.* Crown 8vo. cloth, 10s. 6d.

The Elements of Greek Accentuation (for Schools): abridged from his larger work by H. W. Chandler, M.A., Waynflete Professor of Moral and Metaphysical Philosophy, Oxford. Ext. fcap. 8vo. cloth, 2s. 6d.

A Series of Graduated Greek Readers.

- First Greek Reader.** *In the Press.*
- Second Greek Reader.** *In Preparation.*
- Third Greek Reader.** *In Preparation.*
- Fourth Greek Reader; being Specimens of Greek Dialects.** With Introductions and Notes. By W. W. Merry, M.A., Fellow and Lecturer of Lincoln College. Extra fcap. 8vo. cloth, 4s. 6d.
- Fifth Greek Reader.** Part I. Selections from Greek Epic and Dramatic Poetry, with Introductions and Notes. By Evelyn Abbott, M.A., Fellow of Balliol College. Ext. fcap. 8vo. cloth, 4s. 6d.
- Part II. By the same Editor. *In Preparation.*
- Xenophon. Easy Selections** (for Junior Classes). With a Vocabulary, Notes, and Map. By J. S. Phillpotts, B.C.L., Head Master of Bedford School; and C. S. Jerram, M.A., late Scholar of Trinity College, Oxford. Extra fcap. 8vo. cloth, 3s. 6d.
- Xenophon. Selections** (for Schools). With Notes and Maps. By J. S. Phillpotts, B.C.L., Head Master of Bedford School. Part I. *Third Edition.* Ext. fcap. 8vo. cloth, 3s. 6d.
- Arrian. Selections** (for Schools). With Notes. By J. S. Phillpotts, B.C.L., Head Master of Bedford School.
- The Golden Treasury of Ancient Greek Poetry; being a Collection of the finest passages in the Greek Classic Poets, with Introductory Notices and Notes.** By R. S. Wright, M.A., Fellow of Oriel College, Oxford. Ext. fcap. 8vo. cloth, 8s. 6d.
- A Golden Treasury of Greek Prose, being a collection of the finest passages in the principal Greek Prose Writers, with Introductory Notices and Notes.** By R. S. Wright, M.A., and J. E. L. Shadwell, M.A. Ext. fcap. 8vo. cloth, 4s. 6d.
- Aristotle's Politics.** By W. L. Newman, M.A., Fellow of Balliol College, Oxford.
- Demosthenes and Aeschines.** The Orations of Demosthenes and Aeschines on the Crown. With Introductory Essays and Notes. By G. A. Simcox, M.A., and W. H. Simcox, M.A. 8vo. cloth, 12s.
- Theocritus** (for Schools). With Notes. By H. Kynaston, M.A. (late Snow), Head Master of Cheltenham College. *Second Edition.* Extra fcap. 8vo. cloth, 4s. 6d.
- Homer. Odyssey, Book II.** With Introduction, Notes, and Table of Homeric Forms. By W. W. Merry, M.A., Fellow and Lecturer of Lincoln College, Oxford. Ext. fcap. 8vo. cloth, 1s. 6d.
- Homer. Odyssey, Books I—XII** (for Schools). By the same Editor. *Fifth Edition.* Extra fcap. 8vo. cloth, 4s. 6d.
- Homer. Odyssey, Books XIII—XXIV** (for Schools). By the same Editor. *In Preparation.*

A Homeric Grammar. By D. B. Monro, M.A., Fellow of Oriel College. *In Preparation.*

Homer. Odyssey, Books I–XII. Edited with English Notes, Appendices, etc. By W. W. Merry, M.A., Fellow and Lecturer of Lincoln College, Oxford; and the late James Riddell, M.A., Fellow of Balliol College, Oxford. Demy 8vo. *cloth*, 16s.

Homer. Odyssey, Books XIII–XXIV. By Robinson Ellis, M.A., Fellow of Trinity College, Oxford.

Homer. Iliad. By D. B. Monro, M.A., Fellow and Tutor of Oriel College, Oxford.

Also a small edition for Schools.

Plato. Selections (for Schools). With Notes. By B. Jowett, M.A., Regius Professor of Greek; and J. Purves, M.A., Fellow and Lecturer of Balliol College, Oxford.

Sophocles. The Plays and Fragments. With English Notes and Introductions. By Lewis Campbell, M.A., Professor of Greek, St. Andrews, formerly Fellow of Queen's College, Oxford. 2 vols.

Vol. I. Oedipus Tyrannus. Oedipus Coloneus. Antigone. 8vo. *cloth*, 14s.

Sophocles. The Text of the Seven Plays. By the same Editor. Ext. fcap. 8vo. *cloth*, 4s. 6d.

Sophocles. In Single Plays, with English Notes, &c. By Lewis Campbell, M.A., Professor of Greek, St. Andrews, and Evelyn Abbott, M.A., Fellow of Balliol College, Oxford. Extra fcap. 8vo. *limp*.

Oedipus Rex, Oedipus Coloneus, Antigone, 1s. 9d. each.
Ajax, Electra, Trachiniae, 2s. each.

Sophocles. Oedipus Rex: Dindorf's Text, with Notes by the present Bishop of St. David's Ext. fcap. 8vo. *limp*, 1s. 6d.

III. MENTAL AND MORAL PHILOSOPHY.

The Elements of Deductive Logic, designed mainly for the use of Junior Students in the Universities. By T. Fowler, M.A., Professor of Logic, Oxford. *Fifth Edition*, with a Collection of Examples. Extra fcap. 8vo. *cloth*, 3s. 6d.

The Elements of Inductive Logic, designed mainly for the use of Students in the Universities. By the same Author. *Second Edition*. Extra fcap. 8vo. *cloth*, 6s.

Selections from Berkeley, with an Introduction and Notes. For the use of Students in the Universities. By Alexander Campbell Fraser, LL.D. Crown 8vo. *cloth*, 7s. 6d. *See also* p. 16.

A Manual of Political Economy, for the use of Schools. By J. E. Thorold Rogers, M.A., formerly Professor of Political Economy, Oxford. *Third Edition*. Extra fcap. 8vo. *cloth*, 4s. 6d.

IV. MATHEMATICS, &c.

Figures Made Easy: a first Arithmetic Book. (Introductory to 'The Scholar's Arithmetic.') By Lewis Hensley, M.A., formerly Fellow and Assistant Tutor of Trinity College, Cambridge. Crown 8vo. *cloth*, 6d.

Answers to the Examples in Figures made Easy, together with two thousand additional Examples formed from the Tables in the same, with Answers. By the same Author. Crown 8vo. *cloth*, 1s.

The Scholar's Arithmetic; with Answers to the Examples. By the same Author. Crown 8vo. *cloth*, 4s. 6d.

The Scholar's Algebra. An Introductory work on Algebra. By the same Author. Crown 8vo. *cloth*, 4s. 6d.

Book-keeping. By R. G. C. Hamilton, Financial Assistant Secretary to the Board of Trade, and John Ball (of the Firm of Quilter, Ball, & Co.), Co-Examiners in Book-keeping for the Society of Arts. *New and enlarged Edition.* Extra fcap. 8vo. *limp cloth*, 2s.

A Course of Lectures on Pure Geometry. By Henry J. Stephen Smith, M.A., F.R.S., Fellow of Corpus Christi College, and Savilian Professor of Geometry in the University of Oxford.

Acoustics. By W. F. Donkin, M.A., F.R.S., Savilian Professor of Astronomy, Oxford. Crown 8vo. *cloth*, 7s. 6d.

A Treatise on Electricity and Magnetism. By J. Clerk Maxwell, M.A., F.R.S., Professor of Experimental Physics in the University of Cambridge. 2 vols. 8vo. *cloth*, 1l. 11s. 6d.

An Elementary Treatise on the same subject. By the same Author. *Preparing.*

V. HISTORY.

A Constitutional History of England, in its Origin and Development. By W. Stubbs, M.A., Regius Professor of Modern History in the University of Oxford. Vols. I and II. Crown 8vo. *cloth*, each 12s.

Vol. III. *In the Press.*

Select Charters and other Illustrations of English Constitutional History, from the Earliest Times to the Reign of Edward I. Arranged and Edited by W. Stubbs, M.A., Regius Professor of Modern History in the University of Oxford. *Second Edition.* Crown 8vo. *cloth*, 8s. 6d.

- A History of England**, principally in the Seventeenth Century.
By Leopold Von Ranke. Translated by Resident Members of the University of Oxford, under the superintendence of G. W. Kitchin, M.A., and C. W. Boase, M.A. 6 vols. 8vo. *cloth*, 3l. 3s.
- Genealogical Tables illustrative of Modern History.** By H. B. George, M.A. *Second Edition. Revised and Corrected.* Small 4to. *cloth*, 12s.
- A History of France.** With numerous Maps, Plans, and Tables. By G. W. Kitchin, M.A. In Three Volumes, crown 8vo. *cloth*, 10s. 6d. *each*.
Vol. 1. Down to the Year 1453. Vol. 2. From 1453-1624.
Vol. 3. From 1624-1793.
- A Manual of Ancient History.** By George Rawlinson, M.A., Camden Professor of Ancient History, formerly Fellow of Exeter College, Oxford. Demy 8vo. *cloth*, 14s.
- A History of Germany** and of the Empire, down to the close of the Middle Ages. By J. Bryce, D.C.L., Regius Professor of Civil Law in the University of Oxford.
- A History of British India.** By S. J. Owen, M.A., Reader in Indian Law and History in the University of Oxford.
- A History of Greece.** By E. A. Freeman, M.A., formerly Fellow of Trinity College, Oxford.
- A Selection from the Despatches, Treaties, and other Papers** of the Marquess Wellesley, K.G., during his Government of India; with Appendix, Map of India, and Plans. Edited by S. J. Owen, M.A., Reader in Law and History, and Tutor at Christ Church; formerly Professor of History in the Elphinstone College, Bombay. 8vo. *cloth*, 1l. 4s. *Just Published.*

VI. LAW.

- An Introduction to the Principles of Morals and Legislation.** By Jeremy Bentham. Crown 8vo. *cloth*, 6s. 6d.
- Elements of Law** considered with reference to Principles of General Jurisprudence. By William Markby, M.A., Judge of the High Court of Judicature, Calcutta. *Second Edition, with Supplement.* Crown 8vo. *cloth*, 7s. 6d.
- An Introduction to the History of the Law of Real Property**, with original Authorities. By Kenelm E. Digby, M.A., of Lincoln's Inn, Barrister-at-Law, and formerly Fellow of Corpus Christi College, Oxford. Crown 8vo. *cloth*, 7s. 6d.
- The Elements of Jurisprudence.** By Thomas Erskine Holland, D.C.L., Chichele Professor of International Law and Diplomacy, and formerly Fellow of Exeter College, Oxford. *Preparing.*
- The Institutes of Justinian**, edited as a recension of the Institutes of Gaius. By the same Editor. Extra fcap. 8vo. *cloth*, 5s.

Gaii Institutionum Juris Civilis Commentarii Quatuor; or, Elements of Roman Law by Gaius. With a Translation and Commentary by Edward Poste, M.A., Barrister-at-Law, and Fellow of Oriel College, Oxford. *Second Edition.* 8vo. cloth, 18s.

Select Titles from the Digest of Justinian. By T. E. Holland, D.C.L., Chichele Professor of International Law and Diplomacy, and formerly Fellow of Exeter College, Oxford, and C. L. Shadwell, B.C.L., Fellow of Oriel College, Oxford. *In Parts.*

Part I. Introductory Titles. 8vo. sewed, 2s. 6d.

Part II. Family Law. 8vo. sewed, 1s.

Part III. Property Law. 8vo. sewed, 2s. 6d.

VII. PHYSICAL SCIENCE.

A Handbook of Descriptive Astronomy. By G. F. Chambers, F.R.A.S., Barrister-at-Law. Third Edition. Demy 8vo. cloth, 28s.

Chemistry for Students. By A. W. Williamson, Phil. Doc., F.R.S., Professor of Chemistry, University College, London. *A new Edition, with Solutions.* Extra fcap. 8vo. cloth, 8s. 6d.

A Treatise on Heat, with numerous Woodcuts and Diagrams. By Balfour Stewart, LL.D., F.R.S., Professor of Natural Philosophy in Owens College, Manchester. *Third Edition.* Extra fcap. 8vo. cloth, 7s. 6d.

Forms of Animal Life. By G. Rolleston, M.D., F.R.S., Linacre Professor of Physiology, Oxford. Illustrated by Descriptions and Drawings of Dissections. Demy 8vo. cloth, 16s.

Exercises in Practical Chemistry (Laboratory Practice). By A. G. Vernon Harcourt, M.A., F.R.S., Senior Student of Christ Church, and Lee's Reader in Chemistry; and H. G. Madan, M.A., Fellow of Queen's College, Oxford.

Series I. Qualitative Exercises. *Second Edition.* Crown 8vo. cloth, 7s. 6d.

Series II. Quantitative Exercises.

Geology of Oxford and the Valley of the Thames. By John Phillips, M.A., F.R.S., Professor of Geology, Oxford. 8vo. cloth, 21s.

Text-Book of Botany, Morphological and Physiological. By Dr. Julius Sachs, Professor of Botany in the University of Würzburg. Translated by A. W. Bennett, M.A., Lecturer on Botany, St. Thomas' Hospital, assisted by W. T. Thiselton Dyer, M.A., Christ Church. Royal 8vo. half morocco, 31s. 6d.

Crystallography. By M. H. N. Story-Maskelyne, M.A., Professor of Mineralogy, Oxford; and Deputy Keeper in the Department of Minerals, British Museum. *In the Press.*

VIII. ENGLISH LANGUAGE AND LITERATURE.

A First Reading Book. By Marie Eichens of Berlin; and edited by Anne J. Clough. Extra fcap. 8vo. *stiff covers*, 4d.

Oxford Reading Book, Part I. For Little Children. Extra fcap. 8vo. *stiff covers*, 6d.

Oxford Reading Book, Part II. For Junior Classes. Extra fcap. 8vo. *stiff covers*, 6d.

On the Principles of Grammar. By E. Thring, M.A., Head Master of Uppingham School. Extra fcap. 8vo. *cloth*, 4s. 6d.

Grammatical Analysis, designed to serve as an Exercise and Composition Book in the English Language. By E. Thring, M.A., Head Master of Uppingham School. Extra fcap. 8vo. *cloth*, 3s. 6d.

An English Grammar and Reading Book, for Lower Forms in Classical Schools. By O. W. Tancock, M.A., Assistant Master in Sherborne School. *Second Edition*. Extra fcap. 8vo. *cloth*, 3s. 6d.

The Philology of the English Tongue. By J. Earle, M.A., formerly Fellow of Oriel College, and sometime Professor of Anglo-Saxon, Oxford. *Second Edition*. Extra fcap. 8vo. *cloth*, 7s. 6d.

A Book for the Beginner in Anglo-Saxon, comprising a short Grammar and some Selections from the Gospels. By John Earle, M.A., Professor of Anglo-Saxon in the University of Oxford. Extra fcap. 8vo. *cloth*, 2s. 6d.

An Anglo-Saxon Reader. In Prose and Verse. With Grammatical Introduction, Notes, and Glossary. By Henry Sweet, M.A. Extra fcap. 8vo. *cloth*, 8s. 6d.

Chaucer. The Prioresses Tale; Sir Thopas; The Monkes Tale; The Clerkes Tale; The Squieres Tale, &c. Edited by W. W. Skeat, M.A., Editor of *Piers the Plowman*, &c., &c. Extra fcap. 8vo. *cloth*, 4s. 6d.

Chaucer. The Tale of the Man of Lawe; The Pardoner's Tale; The Second Nonnes Tale; The Chanouns Yemannes Tale. By the same Editor. Extra fcap. 8vo. *cloth*, 4s. 6d. *Just Published*.

Specimens of Early English. A New and Revised Edition. With Introduction, Notes, and Glossarial Index. By R. Morris, LL.D., and W. W. Skeat, M.A.

Part I. *In the Press*.

Part II. From Robert of Gloucester to Gower (A.D. 1298 to A.D. 1393). *Second Edition*. Extra fcap. 8vo. *cloth*, 7s. 6d.

Specimens of English Literature, from the 'Ploughmans Crede' to the 'Shepheardes Calender' (A.D. 1394 to A.D. 1579). With Introduction, Notes, and Glossarial Index. By W. W. Skeat, M.A. Extra fcap. 8vo. *cloth*, 7s. 6d.

The Vision of William concerning Piers the Plowman, by William Langland. Edited, with Notes, by W. W. Skeat, M.A. *Second Edition*. Extra fcap. 8vo. *cloth*, 4s. 6d.

Shakespeare. *Hamlet*. Edited by W. G. Clark, M.A., and W. Aldis Wright, M.A. Extra fcap. 8vo. *stiff covers*, 2s.

Shakespeare. *The Tempest*. By W. Aldis Wright, M.A. Extra fcap. 8vo. *stiff covers*, 1s. 6d.

Shakespeare. *King Lear*. By the same Editor. Extra fcap. 8vo. *stiff covers*, 1s. 6d.

Shakespeare. *As You Like It*. By the same Editor. Extra fcap. 8vo. *stiff covers*, 1s. 6d. (For other Plays, see p. 31.)

Shakespeare. *Midsummer Night's Dream*. By the same Editor. *In the Press*.

Milton. *The Areopagitica*. With Introduction and Notes. By J. W. Hales, M.A., late Fellow of Christ's College, Cambridge. Extra fcap. 8vo. *cloth*, 3s.

Addison. *Selections from Papers in the Spectator*. With Notes. By T. Arnold, M.A., University College. Extra fcap. 8vo. *cloth*, 4s. 6d.

Typical Selections from the best English Writers with Introductory Notices. *Second Edition*. In Two Volumes. Extra fcap. 8vo. *cloth*, 7s. *Sold separately*, price 3s. 6d. each.

Vol. I. Latimer to Berkeley. Vol. II. Pope to Macaulay.

See also XIII. below for other English Classics.

IX. FRENCH LANGUAGE AND LITERATURE.

An Etymological Dictionary of the French Language, with a Preface on the Principles of French Etymology. By A. Brachet. Translated into English by G. W. Kitchin, M.A., formerly Censor of Christ Church. Crown 8vo. *cloth*, 10s. 6d.

Brachet's Historical Grammar of the French Language. Translated into English by G. W. Kitchin, M.A. Third Edition. Extra fcap. 8vo. *cloth*, 3s. 6d.

Corneille's Cinna, and **Molière's Les Femmes Savantes**. Edited, with Introduction and Notes, by Gustave Masson. Extra fcap. 8vo. *cloth*, 2s. 6d.

Racine's Andromaque, and **Corneille's Le Menteur**. With Louis Racine's Life of his Father. By the same Editor. Extra fcap. 8vo. *cloth*, 2s. 6d.

Molière's Les Fourberies de Scapin, and **Racine's Athalie**. With Voltaire's Life of Molière. By the same Editor. Extra fcap. 8vo. *cloth*, 2s. 6d.

Selections from the Correspondence of **Madame de Sévigné** and her chief Contemporaries. Intended more especially for Girls' Schools. By the same Editor. Extra fcap. 8vo. cloth, 3s.

Voyage autour de ma Chambre, by **Xavier de Maistre**; **Ourika**, by **Madame de Duras**; **La Dot de Suzette**, by **Fievée**; **Les Jumeaux de l'Hôtel Corneille**, by **Edmond About**; **Mésaventures d'un Écolier**, by **Rodolphe Töpffer**. By the same Editor. Extra fcap. 8vo. cloth, 2s. 6d.

Regnard's Le Joueur, and **Brueys and Palaprat's Le Grondeur**. By the same Editor. Extra fcap. 8vo. cloth, 2s. 6d.

Louis XIV and his Contemporaries; as described in Extracts from the best Memoirs of the Seventeenth Century. With English Notes, Genealogical Tables, &c. By the same Editor. Extra fcap. 8vo. cloth, 2s. 6d.

X. GERMAN LANGUAGE AND LITERATURE.

New German Method. In Four Volumes. By **Hermann Lange**, Teacher of Modern Languages, Manchester.

Vol. I. **The Germans at Home**. 8vo. cloth, 2s. 6d.

Vol. II. **A Grammar of the German Language**. 8vo. cloth, 3s. 6d.

Vol. III. **The German Manual**. *In the Press*.

Vol. IV. **German Composition**. *In Preparation*.

Lessing's Laocoon. With Introduction, English Notes, etc. By **Dr. Albert Haman**, Teacher of German at the Taylor Institution, Oxford. *In Preparation*.

Also, *Edited by* **C. A. BUCHHEIM**, *Phil. Doc., Professor in King's College, London*.

Goethe's Egmont. With a Life of Goethe, &c. Extra fcap. 8vo. cloth, 3s.

Schiller's Wilhelm Tell. With a Life of Schiller; an historical and critical Introduction, Arguments, and a complete Commentary. *Second Edition*. Extra fcap. 8vo. cloth, 3s. 6d.

Lessing's Minna von Barnhelm. A Comedy. With a Life of Lessing, Critical Analysis, Complete Commentary, &c. *Second Edition*. Extra fcap. 8vo. cloth, 3s. 6d.

In Preparation.

Goethe's Iphigenie auf Tauris. A Drama. With a Critical Introduction, Arguments to the Acts, and a complete Commentary.

Selections from the Poems of Schiller and Goethe.

Becker's (K. F.) Friedrich der Grosse.

Egmont's Leben und Tod, and **Belagerung von Antwerpen** by Schiller.

XI. ART, &c.

- A Handbook of Pictorial Art.** By R. St. J. Tyrwhitt, M.A. formerly Student and Tutor of Christ Church, Oxford. With coloured Illustrations, Photographs, and a chapter on Perspective by A. Macdonald. *Second Edition.* 8vo. *half morocco*, 18s.
- A Music Primer for Schools.** By J. Troutbeck, M.A., Music Master in Westminster School, and R. F. Dale, M.A., B. Mus., Assistant Master in Westminster School. Crown 8vo. *cloth*, 1s. 6d.
- A Treatise on Harmony.** By Sir F. A. Gore Ouseley, Bart., Professor of Music in the University of Oxford. *Second Edition.* 4to. *cloth*, 10s.
- A Treatise on Counterpoint, Canon, and Fugue**, based upon that of Cherubini. By the same Author. 4to. *cloth*, 16s.
- A Treatise on Musical Form and General Composition.** By the same Author. 4to. *cloth*, 10s.
- The Cultivation of the Speaking Voice.** By John Hullah. *Second Edition.* Extra fcap. 8vo. *cloth*, 2s. 6d.

XII. MISCELLANEOUS.

- Dante.** Selections from the *Inferno*. With Introduction and Notes. By H. B. Cotterill, B.A. Extra fcap. 8vo. *cloth*, 4s. 6d.
- Tasso. La Gerusalemme Liberata.** Cantos i, ii. With Introduction and Notes. By the same Editor. Extra fcap. 8vo. *cloth*, 2s. 6d.
- A Treatise on the use of the Tenses in Hebrew.** By S. R. Driver, M.A., Fellow of New College. Extra fcap. 8vo. *cloth*, 6s. 6d.
- Outlines of Textual Criticism applied to the New Testament.** By C. E. Hammond, M.A., Fellow and Tutor of Exeter College, Oxford. *Second Edition.* Extra fcap. 8vo. *cloth*, 3s. 6d.
- Ancient Liturgies.** With Introduction, Notes, &c. By C. E. Hammond, M.A. *In Preparation.*
- The Modern Greek Language** in its relation to Ancient Greek. By E. M. Geldart, B.A. Extra fcap. 8vo. *cloth*, 4s. 6d.
- A System of Physical Education : Theoretical and Practical.** By Archibald Maclaren. Extra fcap. 8vo. *cloth*, 7s. 6d.

XIII. A SERIES OF ENGLISH CLASSICS,

Designed to meet the wants of Students in English Literature, under the superintendence of the Rev. J. S. BREWER, M.A., of Queen's College, Oxford, and Professor of English Literature at King's College, London.

It is also especially hoped that this Series may prove useful to Ladies' Schools and Middle Class Schools; in which English Literature must always be a leading subject of instruction.

A General Introduction to the Series. By Professor Brewer, M.A.

1. **Chaucer.** The Prologue to the Canterbury Tales; The Knights Tale; The Nonne Prestes Tale. Edited by R. Morris, Editor of Specimens of Early English, &c., &c. *Sixth Edition.* Extra fcap. 8vo. cloth, 2s. 6d. (See also p. 27.)
2. **Spenser's Faery Queene.** Books I and II. Designed chiefly for the use of Schools. With Introduction, Notes, and Glossary. By G. W. Kitchin, M.A., formerly Censor of Christ Church.

Book I. *Eighth Edition.* Extra fcap. 8vo. cloth, 2s. 6d.

Book II. *Third Edition.* Extra fcap. 8vo. cloth, 2s. 6d.
3. **Hooker.** Ecclesiastical Polity, Book I. Edited by R. W. Church, M.A., Dean of St. Paul's; formerly Fellow of Oriel College, Oxford. *Second Edition.* Extra fcap. 8vo. cloth, 2s.
4. **Shakespeare.** Select Plays. Edited by W. G. Clark, M.A., Fellow of Trinity College, Cambridge; and W. Aldis Wright, M.A., Trinity College, Cambridge. Extra fcap. 8vo. stiff covers.

I. The Merchant of Venice. 1s.

II. Richard the Second. 1s. 6d.

III. Macbeth. 1s. 6d. (For other Plays, see p. 28.)
5. **Bacon.**

I. Advancement of Learning. Edited by W. Aldis Wright, M.A. *Second Edition.* Extra fcap. 8vo. cloth, 4s. 6d.

II. The Essays. With Introduction and Notes. By J. R. Thursfield, M.A., Fellow and Tutor of Jesus College, Oxford.
6. **Milton.** Poems. Edited by R. C. Browne, M.A., and Associate of King's College, London. 2 vols. *Fourth Edition.* Extra fcap. 8vo. cloth, 6s. 6d.


Sold separately, Vol. I. 4s.; Vol. II. 3s.


7. **Dryden.** *Select Poems.* Stanzas on the Death of Oliver Cromwell; *Astræa Redux*; *Annus Mirabilis*; *Absalom and Achitophel*; *Religio Laici*; *The Hind and the Panther.* Edited by W. D. Christie, M.A., Trinity College, Cambridge. *Second Edition.* Ext. fcap. 8vo. cloth, 3s. 6d.
8. **Bunyan.** *The Pilgrim's Progress; Grace Abounding.* Edited by E. Venables, M.A., Canon of Lincoln. *In the Press.*
9. **Pope.** With Introduction and Notes. By Mark Pattison, B.D., Rector of Lincoln College, Oxford.
 - I. *Essay on Man.* *Fifth Edition.* Extra fcap. 8vo. stiff covers, 1s. 6d.
 - II. *Satires and Epistles.* *Second Edition.* Extra fcap. 8vo. stiff covers, 2s.
10. **Johnson.** *Rasselas; Lives of Pope and Dryden.* By T. Arnold, M.A., University College.
11. **Burke.** *Select Works.* Edited, with Introduction and Notes, by E. J. Payne, M.A., of Lincoln's Inn, Barrister-at-Law, and Fellow of University College, Oxford.
 - I. *Thoughts on the Present Discontents; the two Speeches on America.* *Second Edition.* Extra fcap. 8vo. cloth, 4s. 6d.
 - II. *Reflections on the French Revolution.* Extra fcap. 8vo. cloth, 3s.
12. **Cowper.** Edited, with Life, Introductions, and Notes, by H. T. Griffith, B.A., formerly Scholar of Pembroke College, Oxford.
 - I. *The Didactic Poems of 1782, with Selections from the Minor Pieces, A.D. 1779-1783.* Extra fcap. 8vo. cloth, 3s.
 - II. *The Task, with Tirocinium, and Selections from the Minor Poems, A.D. 1784-1799.* Extra fcap. 8vo. cloth, 3s.


Published for the University by
MACMILLAN AND CO., LONDON.

Also to be had at the
CLARENDON PRESS DEPOSITORY, OXFORD.

The DELEGATES OF THE PRESS invite suggestions and advice from all persons interested in education; and will be thankful for hints, &c. addressed to the SECRETARY TO THE DELEGATES, Clarendon Press, Oxford.







14 DAY USE
RETURN TO DESK FROM WHICH BORROWED

LOAN DEPT.

This book is due on the last date stamped below, or
on the date to which renewed.
Renewed books are subject to immediate recall.

APR 18 1966 3 3

REC'D LD
APR 18 1966

MUN 7 1963

REC. CIR. DEC 09 '82

LD 21A-60m-10,'65
(F7763s10)476B

General Library
University of California
Berkeley

20m-6, 61 (C63954)

YC 40029

14008

QC 175

N 3

